arXiv:hep-th/9806177v2 8 Jul 1998

hep-th/9806177, IASSNS-HEP-98/58

Brane Box Realization of Chiral Gauge Theories in Two Dimensions

Hugo García-Compeán[1], Angel M. Uranga[2],

School of Natural Sciences

Institute for Advanced Study

Olden Lane, Princeton, NJ 08540, USA

Abstract

We study type IIA configurations of D4 branes and three kinds of NS fivebranes. The D4 brane world-volume has finite extent in three directions, giving rise to a two-dimensional low-energy field theory. The models have generically $(0,2)$ supersymmetry. We determine the rules to read off the spectrum and interactions of the field theory from the brane box configuration data. We discuss the construction of theories with enhanced $(0,4)$, $(0,6)$ and $(0,8)$ supersymmetry. Using T-duality along the directions in which the D4 branes are finite, the configuration can be mapped to D1 branes at $\mathbb{C}^4/\Gamma$ singularities, with Γ an abelian subgroup of $SU(4)$. This provides a rederivation of the rules in the brane box model. The enhancement of supersymmetry has a nice geometrical interpretation in the singularity picture in terms of the holonomy group of the four-fold singularity.

June, 1998

[1] Also *Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN, Apdo. Postal 14-740, 07000, México D.F., México* E-mail: compean@sns.ias.edu

[2] E-mail: uranga@sns.ias.edu

1. Introduction

The dynamics of D-branes in certain configurations of intersecting branes encodes many field-theoretical facts about supersymmetric theories in several dimensions (for a review, see [1]). Gauge theories in $p+1$ dimensions with sixteen supercharges can be obtained as the world-volume theories of flat infinite Dp-branes. In the context of theories with eight supersymmetries in p dimensions, it was shown in [2] that such theories can be realized by considering Dp-branes with a world-volume which is finite in one direction, in which the D brane ends on NS fivebranes. For definiteness, let us take such world-volume spanning $0, 1, 2, \ldots, (p-1)$, and with finite extent along 6. The brane is suspended between NS fivebranes spanning 012345. The low energy theory in the non-compact dimensions of the D-brane is p-dimensional. It is still a gauge theory, but the presence of the NS branes breaks half of the supersymmetries, so eight supercharges remain. This construction has been generalized in several directions, and has yielded the realization of a large family of models in several dimensions. This setup has also been exploited to compute different exact quantum results in these theories. For a review of such achievements, see [1].

A nice property of the interplay of field theories and configurations of branes is that the intersections of branes can sometimes support chiral zero modes. This opens the possibility of studying chiral gauge theories using branes. The simplest such example is provided by the realization of six-dimensional theories with eight supersymmetries, which are chiral. These can be realized in the setup described above by taking $p = 6$, *i.e.* one considers D6 branes extending along 0123456, and which are bounded in 6 by NS branes with world-volume along 012345. This construction has been discussed at length in [3], where it was shown that the configurations yield theories satisfying the very restricting anomaly cancellation conditions. The family of models obtained reproduces the results of [4,5], and even contains some further consistent examples.

Chirality if a fragile property, in the sense that toroidal compactifications or too much supersymmetry spoil it. Thus, in order to obtain chiral theories in four dimensions one has to consider theories with only four supercharges. Their realization in terms of branes requires new ingredients. A fairly general family of brane configurations realizing generically chiral gauge theories in four dimensions was introduced in [6][3]. The idea is a

[3] Let us also mention that the approach of rotated branes [7] has been used to construct some chiral gauge theories in four dimensions [8]. This construction, however, seems not so flexible and easy to generalize. An interesting apporach, introduced in [9], is related to the constructions of [6] by T-duality.

clever extension of the philosophy in [2]. It consists in realizing first a five-dimensional theory with eight supercharges, by using D5 branes along 012346, suspended between NS branes with world-volume along 012345. Then, the D5 brane is bounded in the direction 4, by using a new set of NS branes oriented along 012367 (denoted NS$'$ branes). The low energy theory is four-dimensional, since the world-volume of the D5 brane along 46 is a finite rectangle. Such configurations are known as brane box models. The presence of the new kind of branes breaks a further half of the supersymmetries, and so the theory has only four supercharges. Furthermore, the intersections of NS, NS$'$ and D5 branes introduce chirality in the four dimensional theory. There is no complete understanding of the quantum effects of these gauge theories in terms of branes, even though some results on exact finiteness and marginality were obtained in [10]. Some understanding on the bending of the branes in this configurations, recently developed in [11], may help in improving the situation in this respect.

In this paper we continue this analysis of chirality in field theories realized by brane configurations, and construct chiral gauge theories in two dimensions, with two supercharges, *i.e.* $(0,2)$ theories. After the previous discussion, there is a natural approach to the construction of such theories. We first realize three-dimensional gauge theories with four supercharges, by using D4 branes along 01246 and NS, NS$'$ branes as before. The configuration is T-dual (along 3) to the brane box models constructed above. We then bound the D4 branes in the direction 2, by means of a new set of NS fivebranes, along 014567, denoted NS$''$ branes. This implies the the low energy field theory will be two-dimensional, and that only two supersymmetries remain unbroken. Now it is the intersection of D4, NS, NS$'$ and NS$''$ branes that introduces the chirality in the two dimensional theory. The construction and study (at the classical level) of these theories is the aim of the present paper.

We start in Section 2 by reviewing some basic features of two-dimensional field theories. In Section 3 we describe the brane configurations sketched above. The spectrum and interactions corresponding to a given brane model is determined by studying first the realization of $(2,2)$ theories, and generalizing the result.

In Section 4 we discuss models with enhanced supersymmetry. An interesting feature of two-dimensional theories is that it is possible to enhance the supersymmetry while preserving chirality. So it makes sense to ask whether new ingredients are required to realize *e.g.* $(0,4)$ theories using brane constructions. In Section 4 we show a simple way

of realizing a large class of models with such enhanced chiral supersymmetry in our setup, without the need of extra ingredients.

In Section 5 we discuss the relation or our approach to that of [15]. Recently it has become clear that there is an alternative way of realizing chiral theories in several dimensions, as the world-volume theories of D brane probes at singularities. Thus, using D5 branes at ADE singularities (possibly with some orientifold projection) one can realize six-dimensional theories with eight supercharges [13,14,5]. Some of these constructions (corresponding to A_k singularities) have been argued to be T-dual to the brane configurations in [3]. Similarly, by using D3 branes at $\mathbb{C}^3/\Gamma$ singularities, with Γ a discrete subgroup of $SU(3)$, one can realize chiral four-dimensional theories with four supercharges [16,17,18,19]. For abelian discrete groups, these constructions are related by T-duality to the brane box models, as argued in [12]. In Section 5 we apply the same argument to the our brane models, and discuss their relation with the theories on the world-volume of D1 branes at $\mathbb{C}^4/\Gamma$ singularities, with Γ a discrete subgroup of $SU(4)$. These latter theories have been studied in [15], and we review the determination of the spectrum and interactions in the singularity language. This can be compared with the rules proposed in Section 3, and provide a rederivation of the result. Also, the agreement supports our T-duality proposal in this case. Finally, it provides a nice geometrical interpretation for the enhancement to $(0,4)$ and $(0,8)$ supersymmetry introduced in Section 4. The explanation suggests the brane configurations with enhanced supersymmetry have an appropriate 'generalized holonomy group' in the sense of [20].

Finally, Section 6 contains our conclusions. The definition of the gauge theory parameters in terms of branes is discussed in an appendix.

We note that brane configurations have been used to obtain several results on non-chiral theories in two dimensions, in [21] for $(2,2)$ theories, and in [22] for $(4,4)$ theories. Another approach to the study of $(2,2)$ theories has been taken in [23] in the framework of geometric engineering.

2. Overview of Field Theory

In this section we will give an overview of the matter content and interactions of the two-dimensional (2,2) and (0,2) gauge theories. Our aim is not to provide an extensive

review of such theories, but to briefly recall the relevant structure of multiplets and interactions, which we will need in the following sections. For a more complete treatment see Refs. [24,25].

2.1. $\mathcal{N} = 1$ Theories in Four Dimensions

Let us first briefly review the matter content and interactions of $\mathcal{N} = 1$ gauge theories in four dimensions. For definiteness let us take the gauge group to be $U(N)$. The theory is formulated in the superspace consisting of four spacetime coordinates x^μ, $(\mu = 0, 1, 2, 3)$ and odd coordinates θ^α, $\overline{\theta}^{\dot{\alpha}}$ The multiplets used in the construction of gauge theories are:

i)-. The *chiral multiplet*, which we will denote by $\widetilde{\Phi}$, which contains a complex scalar and a chiral fermion, in any representation of the gauge group.

*ii)-.*The *gauge multiplet*, $\widetilde{V}$, containing (in the Wess-Zumino gauge) gauge bosons and a Majorana fermion, both in the adjoint representation of the gauge group.

The interactions of the $\mathcal{N} = 1$ theory are encoded in the superspace Lagrangian

$$L = \int d^4x\, d^4\theta \sum_i \left(\overline{\widetilde{\Phi}}_i \exp\left(\sum\nolimits_a T_a \widetilde{V}_a\right) \widetilde{\Phi}_i \right) + \int d^4x\, d^2\theta\, \mathrm{Tr}(W^\alpha W_\alpha)$$
$$+ \int d^4x\, d^2\theta\; W(\widetilde{\Phi}_i) + h.c. - r \int d^4x\, d^4\theta\; \mathrm{Tr}(\widetilde{V}) \qquad (2.1)$$

where $W_\alpha = -\frac{1}{4}\overline{D}^2 D_\alpha \widetilde{V}$ is the gauge field strength chiral superfield. The first term contains the kinetic term and gauge couplings for the chiral multiplets. Here T_a are the generators of the gauge group in the appropriate representation. The second term is the kinetic energy and gauge interactions for the gauge multiplets. The third term is the superpotential; $W(\widetilde{\Phi}_i)$ is a holomorphic function of the chiral superfields $\widetilde{\Phi}_i$. In the applications we are to consider, the superpotential will be cubic, and contains the Yukawa interactions and scalar potential terms. The last term is the Fayet-Iliopoulos term which arises if there is a U(1) factor in the gauge group. Finally let us mention that $\mathcal{N} = 1$ gauge theories have a U(1) R-symmetry. This symmetry can be broken to a discrete subgroup by instanton effects.

2.2. (2,2) Theories in Two Dimensions

We now describe how dimensional reduction of $\mathcal{N} = 1$ in four dimensions yields (2,2) gauge theories in two dimensions. To this end, one takes all fields to be independent

of x^2, x^3, and decomposes the representations of the four-dimensional Lorentz group with respect to the two-dimensional one. The resulting gauge theory in two dimensions is nicely described in the (2,2) superspace $(y^\alpha, \theta^+, \theta^-, \overline{\theta}^+, \overline{\theta}^-)$, where $(y^0, y^1) = (x^0, x^1)$.

The structure of multiplets is very similar to the four-dimensional one, as follows.

i)-. There is a (2,2) *chiral multiplet* Φ, containing a complex scalar ϕ, and two fermions of opposite chirality ψ_+, ψ_-.

ii)-. There is also a *vector multiplet* V, containing gauge bosons, v_α, $\alpha = 0, 1$, two Majorana fermions λ_+, λ_- and one complex scalar σ (this last arising from the components of the four-dimensional vector along x^2, x^3).

The Lagrangian is obtained by simple reduction of the Lagrangian (2.1). It has the following structure,

$$L = L_{ch} + L_{gauge} + L_W + L_{D,\theta}. \tag{2.2}$$

The term L_{ch} is the kinetic energy and gauge couplings of the chiral superfields Φ

$$\int d^2\, y d^4\theta \sum_i \left(\overline{\Phi}_i exp\big(\sum\nolimits_a T^a V_a\big) \Phi_i \right). \tag{2.3}$$

The term L_{gauge} contains the kinetic energy and gauge interactions for the vector multiplets, and takes the form

$$L_{gauge} = -\frac{1}{g^2} \int d^2y\, d^4\theta \mathrm{Tr}\big(\overline{\Sigma}\Sigma\big), \tag{2.4}$$

where Σ is the $(2,2)$ gauge field strength superfield and g is the gauge coupling constant.

The $(2,2)$ superpotential term L_W is given by

$$L_W = -\int d^2y\, d\theta^+ d\theta^- W(\Phi_i)|_{\overline{\theta}^+ = \overline{\theta}^- = 0} - h.c.\ . \tag{2.5}$$

Finally, the last term contains the Fayet-Iliopoulos and theta angle terms,

$$L_{D,\theta} = \frac{it}{2\sqrt{2}} \int d^2y d\theta^+ d\overline{\theta}^- \mathrm{Tr}\big(\Sigma|_{\theta^- = \overline{\theta}^+ = 0}\big) + h.c. \tag{2.6}$$

where $t = ir + \frac{\theta}{2\pi}$.

Among the symmetries present in (2,2) theories, the R-symmetries play an important role. For these theories there can two U(1) R-symmetries, the right-moving U(1) R-symmetry acts on the right-moving odd coordinates $(\theta^+, \overline{\theta}^+)$ in the form $\theta^+ \to e^{i\beta}\theta^+$, $\overline{\theta}^+ \to e^{-i\beta}\overline{\theta}^+$, leaving θ_- and $\overline{\theta}^-$ invariant. Similar definition holds for the left-moving

U(1) R-symmetry acting on the left-moving $(\theta^-, \overline{\theta}^-)$. Since the fermions are charged under these symmetries they can be anomalous. The condition that the mixed anomaly of the R-symmetries and the $U(1)$ gauge factor in $U(N)$ vanishes is

$$\sum_i Q_i = 0, \tag{2.7}$$

where Q_i is the gauge $U(1)$ charge of the i-th chiral field Φ_i. This ensures the conservation of the left-moving and right-moving R-currents J_L, J_R. This relation also implies the non-renormalization of the coefficient of the Fayet-Iliopoulos term.

2.3. (0,2) Theories in Two Dimensions

We now move on to review the building blocks of (0,2) gauge theories in two dimensions. These theories are described in the (0,2) superspace $(y^\alpha, \theta^+, \overline{\theta}^+)$. There are three basic kinds of multiplets which we will use.

i)-. The $(0,2)$ *gauge multiplet* V', which contains gauge bosons v_α, $\alpha = 0, 1$, and one fermion χ_-.

ii)-. The $(0,2)$ *chiral multiplet* Φ', contains one complex scalar ϕ and one chiral fermion ψ_+.

iii)-. The $(0,2)$ *Fermi multiplet*, Λ, is described by an anticommuting superfield. Its complete θ expansion contains a chiral spinor λ_-, an auxiliary field G, and a holomorphic function E depending on the chiral (0,2) superfields Φ'_i. The Fermi multiplet Λ satisfies the constraint $\overline{\mathcal{D}}_+\Lambda = \sqrt{2}E(\Phi')$, with $\overline{\mathcal{D}}_+E = 0$. Here $\overline{\mathcal{D}}_+$ represents the supersymmetric covariant derivative. The expansion in components for the Fermi superfield is

$$\Lambda = \lambda_- - \sqrt{2}\theta^+ G - i\theta^+\overline{\theta}^+(D_0 + D_1)\lambda_- - \sqrt{2}\overline{\theta}^+ E(\Phi') \tag{2.8}$$

with D_α denoting the usual supersymmetric derivative [4].

Gauge theories involving these fields are described by a Lagrangian with the following structure,

$$L = L_{gauge} + L_{ch} + L_F + L_{D,\theta} + L_J. \tag{2.9}$$

[4] Notice that our definitions of the Fermi multiplets and the function E follow [24], and differ from those in [25].

As usual, L_{gauge} is the kinetic term of the gauge multiplet given by

$$L_{gauge} = \frac{1}{8g^2}\int d^2y d\theta^+ d\overline{\theta}^+ \mathrm{Tr}\big(\overline{\Upsilon}\Upsilon\big) \tag{2.10}$$

where Υ is the field strength of V'.

The term L_{ch} contains the kinetic energy and gauge couplings of the (0,2) chiral superfields Φ'_i. It is given by

$$L_{ch} = -\frac{i}{2}\int d^2y d^2\theta \sum_i \left(\overline{\Phi'_i}(\mathcal{D}_0 - \mathcal{D}_1)\Phi'_i\right), \tag{2.11}$$

where $\mathcal{D}_0$ and $\mathcal{D}_1$ are the (0,2) gauge covariant derivatives with respect to V'.

The term L_F describes the dynamics of the Fermi multiplets Λ, and certain interactions. It is given by

$$L_F = -\frac{1}{2}\int d^2y d^2\theta \sum_a \big(\overline{\Lambda}_a \Lambda_a\big). \tag{2.12}$$

For future convenience, we give the expression of this term in components, as obtained upon substitution of (2.8) in (2.12).

$$L_F \;=\; \int d^2y \sum_a \left\{ i\overline{\lambda}_{-,a}(D_0+D_1)\lambda_{-,a} + |G_a|^2 - |E_a|^2 - \sum_j \big(\overline{\lambda}_{-,a}\frac{\partial E_a}{\partial \phi_j}\,\psi_{+,j} + \frac{\partial \overline{E_a}}{\partial \overline{\phi_j}}\,\overline{\psi}_{+,j}\lambda_{-,a}\big)\right\}. \tag{2.13}$$

The Fayet-Iliopoulos and theta angle terms are encoded in the (0,2) Lagrangian $L_{D,\theta}$ which is written as

$$L_{D,\theta} = \frac{t}{4}\int d^2y d\theta^+ \mathrm{Tr}\big(\Upsilon|_{\overline{\theta}^+=0}\big) + h.c. \tag{2.14}$$

where $t = \frac{\theta}{2\pi} + ir$.

Finally (0,2) models do admit an additional interaction term L_J which depends on a set of holomorphic functions $J^a(\Phi')$ of the chiral superfields. There is one such function for each Fermi superfield. They satisfy the relation $\sum_a E_a J^a = 0$. This interaction is the (0,2) analog of the superpotential, and its Lagrangian L_J is given by

$$L_J = -\frac{1}{\sqrt{2}}\int d^2y d\theta^+ \sum_a \left(\Lambda_a J^a|_{\overline{\theta}^+=0}\right) - h.c. \quad . \tag{2.15}$$

The expansion of this term in components is

$$L_J = -\int d^2y \sum_a \left(G_a J^a + \sum_j \lambda_{-,a}\psi_{+,j}\frac{\partial J^a}{\partial \phi_j}\right) - h.c. \quad . \tag{2.16}$$

After combining the Lagrangians L_F and L_J and solving for the equations of motion for the auxiliary fields G, the relevant interaction terms in the Lagrangian (we are not listing the gauge interactions and D-terms here) are

$$\sum_a \left(|J^a(\phi)|^2 + |E^a(\phi)|^2\right) - \sum_{a,j}\left(\overline{\lambda}_{-,a}\frac{\partial E_a}{\partial \phi_j}\psi_{+,j} + \lambda_a \frac{\partial J^a}{\partial \phi_j}\psi_{+,j} + h.c.\right). \tag{2.17}$$

The first term contains the scalar potential, and the second the Yukawa couplings. Notice that the choice of the functions E and J completely defines the interactions of the theory.

For (0,2) theories in two dimensions we have just one U(1) R-symmetry group acting on the superspace coordinates $(\theta^+, \overline{\theta}^+)$. This is right-moving R-symmetry and it acts as $\theta^+ \to e^{i\beta}\theta^+$, $\overline{\theta}^+ \to e^{-i\beta}\overline{\theta}^+$, leaving θ^- , $\overline{\theta}^-$ invariant. Again, Eq. (2.7) provides the condition to have cancellation of the mixed anomalies.

2.4. Decomposition of (2,2) Superfields In Terms of (0,2) Multiplets

For future convenience it will be useful to decompose the $(2,2)$ theories in terms of (0,2) superfields. Looking at the field content of the multiplets, we conclude that the $(2,2)$ vector multiplet V decomposes as a (0,2) gauge multiplet V' (containing the gauge boson v_α $(\alpha = 0,1)$ and the negative chirality spinor χ_-) and a (0,2) chiral multiplet $\Sigma' = \Sigma|_{\theta^- = \overline{\theta}^- = 0}$ consisting of an scalar σ and the positive chirality spinor χ_+ in the adjoint representation of the gauge group.

As for the (2,2) chiral multiplet Φ_i, it decomposes as a (0,2) chiral multiplet $\Phi_i' = \Phi_i|_{\theta^- = \overline{\theta}^- = 0}$ (containing the complex scalar and ϕ_i and the positive chirality spinor $\psi_{+,i}$) and a (0,2) Fermi multiplet Λ_i (containing the negative chirality spinor $\psi_{-,i}$, denoted $\lambda_{-,i}$ in what follows in order to be consistent with our $(0,2)$ conventions).

This Fermi multiplet is given by $\Lambda_i = \frac{1}{\sqrt{2}}\mathcal{D}_-\Phi_i|_{\theta^- = \overline{\theta}^- = 0}$. It can be verified that the corresponding function E_i of the $(0,2)$ chiral superfields is $E_i = i\sqrt{2}T_a\Sigma_a'\Phi_i'$, where a runs over the generators of the gauge group under which the field is charged, and T_a are in the appropriate representation.

Some of the interactions in the $(0,2)$ theory are basically specified by the gauge group and the representations of the chiral and Fermi multiplets. Only the L_J term and the interactions coming from L_F deserve special discussion. The Lagrangian L_J is obtained by reduction of the $(2,2)$ superpotential, and thus involves the chiral and Fermi fields coming from the $(2,2)$ chiral fields. The Lagrangian takes the the form (2.15) with a specific form for the functions J^i, namely

$$J^i = \frac{\partial W}{\partial \Phi'_i} \tag{2.18}$$

with W being the superpotential of the (2,2) theory. The equation $\sum_i E_i J^i = 0$ follows from gauge invariance of W.

There are also interactions between the $(0,2)$ chiral multiplet Σ' and the chiral and Fermi multiplets Φ'_i, Λ_i. These couplings are gauge interactions from the $(2,2)$ point of view, but in the $(0,2)$ theory appear from the expansion of L_F. They are obtained by substituting in Eq.(2.13) the functions E_i found above.

3. The Brane Configurations

In this section we introduce certain supersymmetric configurations of NS, NS$'$, and NS$''$ branes, and D4 branes in Type IIA superstring theory. They give rise to two-dimensional $(0,2)$ field theories. These configurations are obtained in the spirit of the brane box configurations in [6], by considering D-branes which are finite in several directions. As explained in the introduction, they belong to a natural sequence of brane box models yielding chiral theories in six, four and two dimensions (taking D branes compact in one, two and three directions, respectively).

3.1. Description of the Brane Configurations

Let us consider the ingredients of the brane configurations which we will use in this paper. Brane configurations consist of:

a)-. NS fivebranes located along (012345).

b)-. NS' fivebranes located along (012367).

c)-. NS" fivebranes located along (014567).

d)-. D4 branes located along (01246).

In this configuration the D4 branes are finite in the directions 246. They are bounded in the direction 2 by the NS$''$ branes, in the direction 4 by the NS$'$ branes, and in the direction 6 by the NS branes. For the D4 branes to be suspended in this way, it is necessary that the coordinates of all branes in 89 should be equal. It is also required that two NS branes joined by a D4 brane should have the same position in 7, and analogously that two NS$'$ branes joined by a D4 brane should have the same position in 5, and that two NS$''$ branes should have the same position in 3. In Figure 1 we show the 246 three-dimensional space and illustrate how the NS, NS$'$, and NS$''$ brane bound the D4 brane in three directions.

The low-energy effective field theory on the D4 branes is two-dimensional, since 01 are the only non-compact directions in their world-volume. The presence of each kind of NS fivebrane breaks one half of the supersymmetries, and altogether they break to 1/8 of the original supersymmetry. A further half is broken by the D4 branes, and the world-volume theory has $(0, 2)$ supersymmetry in two dimensions. Since the D brane is bounded by NS fivebranes, the world-volume gauge bosons will not be projected out and there will be a gauge group for each box in the model. The $U(1)_R$ R-symmetry of the field theory is manifest as the rotational symmetry in the directions 89.

We note that there are a variety of other objects that can be introduced in the configuration without breaking the supersymmetry. For instance, there are three kinds of D6 branes that can be introduced, namely D6 branes along 0124789, D6$'$ branes along 0125689, and D6$''$ branes along 0134689. They provide vector-like flavours for the gauge groups. These extensions are quite well-known from other contexts, and we will not study them in the present paper.



Fig. 1. A three-dimensional box with 2 NS branes, 2 NS$'$ branes and 2 NS$''$ branes. The D4 brane world-volume fills the cube bounded by the NS fivebranes. For clarity we have shown only a finite interval of the NS, NS$'$ and NS$''$ branes, which actually extend to infinity along 24, 26 and 46, respectively.

There is a first rough classification we can make in these brane configurations, according to whether the directions 246 are taken compact or not. If some of these directions are non-compact, then there will be some semi-infinite box, which will represent some global symmetry. For definiteness we will center on the case in which all three directions are compact, with lengths R_2, R_4 and R_6. Extension of our results to other cases is straightforward.

A generic configuration consists of a three-dimensional grid of k NS branes, k' NS' branes and k'' NS" branes dividing the 246 torus into a set of $kk'k''$ boxes. We will often think about these configurations as infinite periodic arrays of boxes in $\mathbb{R}^3$, quotiented by an infinite discrete group of translations in a three-dimensional lattice Λ. This point of view is particularly useful to define models in which the unit cell has non-trivial identifications of sides [10,12]. In Section 4 we will present some examples of this last case.

3.2. Motivating the Brane Box Rules

The next question we would like to address is what is the spectrum of the $(0,2)$ gauge theory corresponding to a given brane box configuration. The strategy we are to follow in this section is to first consider a particular family of brane box models, namely those with $k''=1$. Notice that if there are no NS$''$ present, the brane configurations can be thought of as a compactification of the brane box models in [6,10], up to a T-duality along direction 3. The corresponding two-dimensional field theory will be a dimensional reduction of the four-dimensional $\mathcal{N}=1$ gauge theories in [6], thus a $(2,2)$ theory. The spectrum and

interactions of such theory are known from the four-dimensional analysis, where it was shown that fields transforming in bi-fundamental representations could be represented as arrows joining the D-branes in different boxes (representing the open strings stretched between the D branes), and superpotential interactions corresponded to closed triangles of arrows (representing open string interactions). Our purpose is to reinterpret such fields and interactions in the $(0,2)$ language, and find some rules yielding the correct spectrum. These will turn out to have a natural generalization to other brane configurations with several NS$''$ branes, and which yield genuine $(0,2)$ field theories.

Let us mention that the introduction of a *single* NS$''$ brane does not change the theory, so it still has $(2,2)$ supersymmetry. It will be useful to introduce such brane, because then the matter multiplets will appear in a suggestive pattern, easy to generalize to $(0,2)$ models.

So let us consider a $k \times k' \times 1$ box model, with trivial identifications of the sides of the unit cell, such as that depicted in Figure 2 for the particular case $k = 3$, $k' = 4$. Let us also place $n_{a,b}$ D4 branes in the box in the position (a,b) in the grid.



Fig. 2. A $3 \times 4 \times 1$ box model, with its twelve different boxes and its unit cell. The sides of the unit cell are identified to make the direction 246 compact. Equivalently, the brane configuration can be described as an infinite periodic array of unit cells. For clarity we only show one slice of the infinite periodic array along 2. Notice that, in this and the following figures, the numbers in the boxes are labels to distinguish them and do not denote the number of D4 branes in the box.

It will be convenient to imagine the brane configuration as an infinite periodic array of boxes extending in 246. Thus, each box will be labeled by three indices (a,b,c) corresponding to its position in the grid in 6, 4 and 2, respectively. Even though things will not depend on the actual value of c (since the periodicity in 2 is of one box), again it will

be useful to maintain such notation, aiming to a further generalization to be completed in following sections.

Using the $(2,2)$ reduction of the rules in [6], we know that the box (a,b) has vector multiplets giving a $U(n_{a,b})$ gauge group[5]. There are also $(2,2)$ chiral multiplets which corresponded to horizontal, vertical and diagonal arrows in the four-dimensional construction. We denote these fields by ${\Phi^H}_{a,b}$ (in the $(\square,\overline{\square})$ of $U(n_{a,b})\times U(n_{a+1,b})$), ${\Phi^V}_{a,b}$ (in the $(\square,\overline{\square})$ of $U(n_{a,b})\times U(n_{a,b+1})$), and $\Phi^D_{a,b}$ (in the $(\square,\overline{\square})$ of $U(n_{a,b})\times U(n_{a-1,b-1})$). There are three such fields for each box in the model.

Let us decompose this field content with respect to $(0,2)$ multiplets and try to define some arrows in the brane box diagram to represent such fields. As reviewed in section 2.4, the $(2,2)$ vector multiplet at each box gives a $(0,2)$ vector multiplet and a $(0,2)$ chiral multiplet in the adjoint of $U(n_{a,b})$. We can represent this field by an arrow which starts in the box (a,b,c), extends along 2, and ends in another copy of the box $(a,b,c+1)$. This transforms in the adjoint, since in this configuration the boxes that differ only in c are identified due to the periodicity in 2. We will denote this $(0,2)$ field by $N_{a,b,c}$.

The $(2,2)$ chiral field $\Phi^H_{a,b}$ gives rise to a $(0,2)$ chiral field and a $(0,2)$ Fermi multiplet transforming in the $(\square,\overline{\square})$ of $U(n_{a,b})\times U(n_{a+1,b})$. Our proposal is that the chiral multiplet, denoted $H_{a,b,c}$, should be represented by an arrow which goes from the box (a,b,c) to the box $(a+1,b,c)$. The Fermi multiplet, denoted $\Lambda^{(1)}_{a,b,c}$, is however represented by an arrow starting from the box (a,b,c) and ending on the box $(a+1,b,c+1)$. It will be useful to also consider an arrow starting from the box $(a+1,b,c+1)$ and ending on the box (a,b,c), which represents the conjugate of the Fermi multiplet, denoted $\Lambda^{(\overline{1})}_{a+1,b,c+1}$. Notice however that this is not an independent field [6]. Notice how a four-dimensional field splits in two due to the presence of the new kind of NS fivebrane. This is somewhat analogous to how four-dimensional $\mathcal{N}=2$ theories are constructed using brane box models [10].

The $(2,2)$ chiral multiplet $\Phi^V_{a,b}$ analogously yields one $(0,2)$ chiral multiplet and one Fermi multiplet transforming in the $(\square,\overline{\square})$ of $U(n_{a,b})\times U(n_{a,b+1})$. We can represent the chiral multiplet, denoted $V_{a,b,c}$ by an arrow going from the box (a,b,c) to the box $(a,b+1,c)$. The Fermi field denoted $\Lambda^{(2)}_{a,b,c}$ corresponds to an arrow from the box (a,b,c) to

[5] In two dimensions we expect the $U(1)$ factors to be dynamical. The freezing argument in [26] applies to theories in four or more dimensions.

[6] Even though the notation may appear confusing, we stick to denoting the fields by the box their arrow starts at.

the box $(a, b+1, c+1)$. Again we introduce another arrow, going from the box $(a, b+1, c+1)$ to the box (a, b, c), which represents the conjugate $\Lambda^{(\overline{2})}_{a,b+1,c+1}$ of the Fermi multiplet.

Finally, the $(2,2)$ chiral multiplet $\Phi^D_{a,b}$ gives rise to one $(0,2)$ chiral multiplet and one Fermi multiplet transforming in the $(\square, \overline{\square})$ of $U(n_{a,b}) \times U(n_{a-1,b-1})$. The chiral multiplet, denoted $D_{a,b,c}$, is represented by an arrow going from the box (a, b, c) to the box $(a-1, b-1, c-1)$. The Fermi field, denoted $\Lambda^{(3)}_{a,b,c}$, corresponds to an arrow from the box (a, b, c) to the box $(a-1, b-1, c)$. Again, the opposite arrow is taken to represent the conjugate $\Lambda^{(\overline{3})}_{a-1,b-1,c}$.

The arrows defining the $(0,2)$ chiral fields are shown in Figure 4, and those for Fermi multiplets (and their conjugates) in Figure 5. The list of $(0,2)$ multiplets is given in Table 1, along with their origin in the $(2,2)$ theory.



Fig. 3. The vertices of the cube with dashed contour are located at the centers of the cube drawn with continuous lines.



Fig. 4. Arrows representing the (0,2) chiral multiplets. For clarity we show the fields $H_{a,b,c}$, $V_{a,b,c}$ and $N_{a,b,c}$, and the field $D_{a+1,b+1,c+1}$ in separate pictures. In this and following pictures, the cube with dashed contour is as defined in Figure 3.

$(\mathbf{0},\mathbf{2})$ **Field**	**Representation**	$(\mathbf{2},\mathbf{2})$ **Field**
$H_{a,b,c}$	$(\Box_{a,b,c}, \overline{\Box}_{a+1,b,c})$	$\Phi^H_{a,b}$
$\Lambda^{(1)}_{a,b,c}$	$(\Box_{a,b,c}, \overline{\Box}_{a+1,b,c+1})$	
$V_{a,b,c}$	$(\Box_{a,b,c}, \overline{\Box}_{a,b+1,c})$	$\Phi^V_{a,b}$
$\Lambda^{(2)}_{a,b,c}$	$(\Box_{a,b,c}, \overline{\Box}_{a,b+1,c+1})$	
$D_{a,b,c}$	$(\Box_{a,b,c}, \overline{\Box}_{a-1,b-1,c-1})$	$\Phi^D_{a,b}$
$\Lambda^{(3)}_{a,b,c}$	$(\Box_{a,b,c}, \overline{\Box}_{a-1,b-1,c})$	
$N_{a,b,c}$	$(\Box_{a,b,c}, \overline{\Box}_{a,b,c+1})$ (Adj.)	Vector

Table 1: Fields in a generic (2,2) model obtained by reduction of a four-dimensional $\mathcal{N}=1$ model. The matter content is split in (0,2) chiral and Fermi multiplets. The origin of the fields in the (2,2) theory is specified in the last column.

Since the boxes are repeated periodically along 2 with period of one box, there is some ambiguity in the assignation of arrows. Now we show that with the above choice the $(0,2)$ interactions arise from closed triangles of arrows.

For each box there are six closed triangles involving the arrows corresponding to the fields H, V, D and $\Lambda^{(i)}$, $i=1,2,3$. They are shown in figures 6C, 6D and 6E (The meaning of the other triangles will be explained later). Recall that the $(0,2)$ superpotential interactions involve one Fermi multiplet and two chiral multiplets, a feature correctly reproduced by our choice of arrows. As mentioned in section 2.4, the $(0,2)$ interactions in L_J arise from the reduction of the $(2,2)$ superpotential. Specifically, we expect an interaction of type (2.15) involving the $(0,2)$ chiral and Fermi fields arising from $(2,2)$ chiral multiplets, namely $H_{a,b,c}$, $V_{a,b,c}$ and $D_{a,b,c}$, and $\Lambda^{(i)}_{a,b,c}$ $(i=1,2,3)$. The functions J in that equation can be obtained from the superpotential through Eq. (2.18). It is easy to check that these are given by the mentioned triangles. Explicitly, we obtain the following J functions:

$$\begin{aligned} J^{(1)}_{a,b,c} &= V_{a+1,b,c+1}D_{a+1,b+1,c+1} - D_{a+1,b,c+1}V_{a,b-1,c} \\ J^{(2)}_{a,b,c} &= D_{a,b+1,c+1}H_{a-1,b,c} - H_{a,b+1,c+1}D_{a+1,b+1,c+1} \\ J^{(3)}_{a+1,b+1,c} &= H_{a,b,c}V_{a+1,b,c} - V_{a,b,c}H_{a,b+1,c}. \end{aligned} \tag{3.1}$$

The triangles can be thought of as describing the Yukawa interactions that arise from the $(0,2)$ superpotential, as in (2.16).

The remaining triangles contain the information about the $(0,2)$ interaction terms coming from the Lagrangian L_F, as in (2.13). These are defined by the functions E associated to the Fermi multiplets. These can be obtained from the $(2,2)$ gauge interactions. The resulting functions are

$$\begin{array}{rcccl} E^{(1)}_{a,b,c} & = & N_{a,b,c}H_{a,b,c+1} & - & H_{a,b,c}N_{a+1,b,c} \\ E^{(2)}_{a,b,c} & = & N_{a,b,c}V_{a,b,c+1} & - & V_{a,b,c}N_{a,b+1,c} \\ E^{(3)}_{a+1,b+1,c+1} & = & N_{a+1,b+1,c}D_{a+1,b+1,c+1} & - & D_{a+1,b+1,c}N_{a,b,c-1}. \end{array} \tag{3.2}$$

These can be read from the triangles in figures 6A, 6B and 6F, respectively. The triangles can be thought of as describing the Yukawa interactions in (2.13).

These functions satisfy the relation

$$\sum_i \sum_{a,b,c} E^{(i)}_{a,b,c} J^{(i)}_{a,b,c} \,=\, 0. \tag{3.3}$$



Fig. 5. The arrows associated to (0,2) Fermi multiplets and their conjugates. The fields represented correspond to $\Lambda^{(1)}_{a,b,c}$, $\Lambda^{(\overline{1})}_{a+1,b,c+1}$ (figure A) $\Lambda^{(2)}_{a,b,c}$, $\Lambda^{(\overline{2})}_{a,b+1,c+1}$ (figure B), and $\Lambda^{(3)}_{a+1,b+1,c}$, $\Lambda^{(\overline{3})}_{a,b,c}$ (figure C).

Observe the analogy between both types of functions and in the interactions they give rise to. This is the main motivation to introduce the same diagrammatic representation for both. Actually, at the level of component fields, the interactions are very similar. It is only the choice of a specific chiral superfield to appear in the functions E of the Fermi

multiplets (in this case, the fields N) that introduces the difference in the origin of the terms, as some arising from the superpotential, and others from the Lagrangian L_F.

We stress that the same theory could be rewritten in $(0,2)$ superspace with a different choice for this special field. In that case, there would be some re-shuffling of interactions, and terms which originate in the $(0,2)$ superpotential for one choice can appear from L_F for another choice. In the case we have studied, it was natural to take the fields N as special, since they were in the adjoint, and the $(0,2)$ superspace Lagrangian obtained with this choice can be further written in $(2,2)$ superspace. For more general $(0,2)$ theories, to be studied in next section, there is no canonical choice of special field.

3.3. General Rules to Obtain the Two-Dimensional Field Theory

In this section we consider a more general model, with a unit cell formed by $k \times k' \times k''$ boxes. In each box we can place an arbitrary number of D4 branes, denoted $n_{a,b,c}$ (with $a = 1, 2, \ldots, k$, $b = 1, 2, \ldots, k'$ and $c = 1, 2, \ldots, k''$) for the box labeled (a, b, c). Indices a, b and c correspond to positions along the directions 6,4 and 2 respectively. Notice that as we define the theory on the torus $\mathbf{T}^3$ the indices a, b, c are defined modulo k, k', k'' respectively.

The rules we determined in the previous section have a natural generalization to this more general case. In the following we state these general rules, and then present some arguments supporting it.

The gauge group and matter content are specified by k, k', k'' and the set of numbers $\{n_{a,b,c}\}$. The gauge group associated to this configuration is given by $\prod_{a,b,c} U(n_{a,b,c})$.

The matter content of the model consists of the following $(0,2)$ chiral multiplets.

- The 'horizontal' multiplet $H_{a,b,c}$, which transforms in the bi-fundamental representation $(\square, \overline{\square})$ of $U(n_{a,b,c}) \times U(n_{a+1,b,c})$. It is represented by an arrow going from the box (a, b, c) to the box $(a+1, b, c)$.
- The 'vertical' multiplet $V_{a,b,c}$, transforming in the $(\square, \overline{\square})$ of $U(n_{a,b,c}) \times U(n_{a,b+1,c})$. It corresponds to an arrow going from the box (a, b, c) to the box $(a, b+1, c)$.
- The 'normal' multiplet $N_{a,b,c}$, in the $(\square, \overline{\square})$ of $U(n_{a,b,c}) \times U(n_{a,b,c+1})$. Its arrow, from the box (a, b, c) to the box $(a, b, c+1)$, is normal to those of H and V.
- The 'diagonal' multiplet $D_{a,b,c}$, which transforms in the $(\square, \overline{\square})$ of $U(n_{a,b,c}) \times U(n_{a-1,b-1,c-1})$. Its arrow goes from the box (a, b, c) to the box $(a-1, b-1, c-1)$.

The arrows corresponding to these chiral fields are depicted in Figure 4.

The theory also contains the following $(0,2)$ Fermi multiplets.

- The conjugate fields $\Lambda^{(1)}_{a,b,c}$, $\Lambda^{(\overline{1})}_{a+1,b,c+1}$, which transform respectively in the $(\square,\overline{\square})$ and $(\overline{\square},\square)$ representations of $U(n_{a,b,c})\times U(n_{a+1,b,c+1})$. They are represented by arrows going from the box (a,b,c) to the box $(a+1,b,c+1)$, and vice versa.
- The conjugate fields $\Lambda^{(2)}_{a,b,c}$, $\Lambda^{(\overline{2})}_{a,b+1,c+1}$, transforming respectively in the $(\square,\overline{\square})$ and $(\overline{\square},\square)$ of $U(n_{a,b,c})\times U(n_{a,b+1,c+1})$. They are represented by arrows from the box (a,b,c) to the box $(a,b+1,c+1)$ and vice versa.
- The conjugate fields $\Lambda^{(3)}_{a,b,c}$, $\Lambda^{(\overline{3})}_{a-1,b-1,c}$, transforming respectively in the $(\square,\overline{\square})$ and $(\overline{\square},\square)$ of $U(n_{a,b,c})\times U(n_{a-1,b-1,c})$. They are represented by arrows from the box (a,b,c) to the box $(a-1,b-1,c)$ and vice versa.

The arrows representing the Fermi supermultiplets and their conjugates are shown in Figure 5. The complete spectrum of matter multiplets is shown in Table 2.

Field	**Representation**
$H_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a+1,b,c})$
$V_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a,b+1,c})$
$N_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a,b,c+1})$
$D_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a-1,b-1,c-1})$
$\Lambda^{(1)}_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a+1,b,c+1})$
$\Lambda^{(\overline{1})}_{a+1,b,c+1}$	$(\square_{a+1,b,c+1},\overline{\square}_{a,b,c})$
$\Lambda^{(2)}_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a,b+1,c+1})$
$\Lambda^{(\overline{2})}_{a,b+1,c+1}$	$(\square_{a,b+1,c+1},\overline{\square}_{a,b,c})$
$\Lambda^{(3)}_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a-1,b-1,c})$
$\Lambda^{(\overline{3})}_{a-1,b-1,c}$	$(\square_{a-1,b-1,c},\overline{\square}_{a,b,c})$

Table 2: Spectrum of chiral matter obtained from the general rules. The table shows the representation of the gauge group in which each field is transforming. We have listed the Fermi multiplets along with their conjugates.

We again stress that there are only three independent Fermi multiplets, the others being merely conjugates of them. However there is no canonical choice of 'fundamental' and 'conjugate' fields. This ambiguity is actually related to the choice of which interactions arise from the $(0,2)$ superpotential from L_J and which from the Lagrangian L_F. This splitting of the interactions is determined by the choice of a 'special' chiral multiplet,

which will appear in all the functions E associated to the Fermi multiplets. The rules below can provide the $(0,2)$ Lagrangian for any choice, but for the ease of explanation we will write the explicit equations only for the case the fields $D_{a,b,c}$ are taken as the special ones. In order to clarify this issue and to complete our set of rules, let us turn to the interactions in the theory.

They are given by closed oriented triangles of arrows in the brane diagram. The possible triangles are depicted in Figure 6. Once a special chiral multiplet is chosen, the interactions are arranged in two sets, those arising from the $(0,2)$ superpotential, and those from the E functions associated to the Fermi multiplets. Since the special field appear in all the E functions, the triangles in which it appears give interactions coming from L_F; below we give rules to compute the corresponding functions E. The triangles where the special field does not appear provide the $(0,2)$ superpotential. We will give rules to compute the corresponding functions J. Since the superpotential (2.18) is holomorphic, we can read from the triangles whose fields Λ are the Fermi multiplets and which are the conjugates.



Fig. 6. Prescriptions to compute the interactions in the (0,2) theory. The twelve triangles depicted encode the definition of the functions E and J corresponding to the Fermi multiplets.

Let us choose the fields $D_{a,b,c}$ to be the special fields. This means that, in figure 6, the triangles 6A, 6B and 6C provide the $(0,2)$ superpotential. It also means that the Fermi superfields are $\Lambda^{(\overline{1})}$, $\Lambda^{(\overline{2})}$ and $\Lambda^{(3)}$, whereas $\Lambda^{(1)}$, $\Lambda^{(2)}$ and $\Lambda^{(\overline{3})}$ are their conjugates. We can read the corresponding functions J from the triangles.

$$\begin{array}{lcll} \Lambda^{(\overline{1})}_{a+1,b,c+1} & \to & J^{(\overline{1})}_{a+1,b,c+1} & = H_{a,b,c}N_{a+1,b,c} - N_{a,b,c}H_{a,b,c+1} \\ \Lambda^{(\overline{2})}_{a,b+1,c+1} & \to & J^{(\overline{2})}_{a,b+1,c+1} & = N_{a,b,c}V_{a,b,c+1} - V_{a,b,c}N_{a,b+1,c} \\ \Lambda^{(3)}_{a+1,b+1,c} & \to & J^{(1)}_{a+1,b+1,c} & = V_{a,b,c}H_{a,b+1,c} - H_{a,b,c}V_{a+1,b,c}. \end{array} \tag{3.4}$$

The remaining triangles, 6D, 6E and 6F, involve the fields D, and so give interactions arising from L_F. The triangles provide the E functions corresponding to the Fermi fields:

$$\begin{array}{lcll} \Lambda^{(\overline{1})}_{a+1,b,c+1} & \to & E^{(\overline{1})}_{a+1,b,c+1} & = D_{a+1,b,c+1}V_{a,b-1,c} - V_{a+1,b,c+1}D_{a+1,b+1,c+1} \\ \Lambda^{(\overline{2})}_{a,b+1,c+1} & \to & E^{(\overline{2})}_{a,b+1,c+1} & = D_{a,b+1,c+1}H_{a-1,b,c} - H_{a,b+1,c+1}D_{a+1,b+1,c+1} \\ \Lambda^{(3)}_{a+1,b+1,c} & \to & E^{(3)}_{a+1,b+1,c} & = D_{a+1,b+1,c}N_{a,b,c-1} - N_{a+1,b+1,c}D_{a+1,b+1,c+1}. \end{array} \tag{3.5}$$

One can check that they verify the relation

$$\sum_i \sum_{a,b,c} \left(J^{(\overline{1})}_{a,b,c}E^{(\overline{1})}_{a,b,c} + J^{(\overline{2})}_{a,b,c}E^{(\overline{2})}_{a,b,c} + J^{(3)}_{a,b,c}E^{(3)}_{a,b,c} \right) = 0. \tag{3.6}$$

Since the functions J and E specify the interactions, we have completed the characterization of the $(0,2)$ field theory. It is a straightforward matter to extract the interactions in components by using the relevant formulae in Section 2. The characterization of the field theory parameters in terms of the parameters in the brane configuration is carried out in appendix I.

This set of rules is the natural generalization of those determined for the particular case of $(2,2)$ theories in Section 3.2. A first good property of these rules is that whenever one of the chiral multiplets appears in adjoint representations, the spectrum and interactions will have enhanced $(2,2)$ supersymmetry. The simplest way of showing it is to take that chiral multiplet as the 'special' one. Then all interactions arising from functions E become gauge interactions in the $(2,2)$ theory, and the couplings form the functions J will correspond to the $(2,2)$ superpotential. We will come back to this and related points in Section 4.

Another check of our set of rules comes from the study of Higgs breakings in the classical theory. It is clear in the brane configuration that there are some brane movements which can be performed. Consider for instance the case with the same number of D4 branes in all boxes. Then the NS branes can be moved along 7, the NS$'$ branes can be moved along 5, and the NS$''$ branes can be moved along 3. As we argue in the appendix, these movements correspond to changing the Fayet-Illiopoulos parameters in the field theory. This, on the other hand, implies that some field in the bi-fundamental gets a vev in order

to make the D-term vanish and maintain unbroken supersymmetry. This triggers gauge symmetry breaking, and as can be checked in the field theory we have proposed, typically pairs of group factors break to the diagonal subgroup. This is precisely the phenomenon observed in the brane picture, and provides support to our identification of the field theory. The analysis is similar to that in [6] and we will not repeat the exercise here.

Another type of Higgs breaking corresponds to recombining D4 branes in different boxes until they complete a set that can separate from the grid of NS fivebranes. For instance, if all boxes have equal number of D4 branes, there is a Higgs branch in which the recombined D4 branes wrap the three-torus completely, and move freely in 35789. Other branches are discussed in analogy with [12]. Of course, our discussion has been in terms of classical language. In two dimensions there is no moduli space of vacua, and our 'Higgs branches' should be understood as the target spaces of the two-dimensional field theory interpreted as a sigma model. The nature of this target space will be further clarified in Section 5.

A related point that usually arises in the context of $(2,2)$ and $(0,2)$ theories is the dependence of the Higgs branch with the FI parameters. In particular, we could ask whether our models will present phase transitions of the kind analyzed in [24], establising some kind of Calabi-Yau/Landau-Ginzburg correspondence. The field theories are rather complicated, and this study is far from straightforward. In Section 5 we will clarify this issue by showing the relation of our field theories to those arising from D-branes at singularities. This allows us to apply several results about the phase structure of these linear sigma models [16,15].

Finally, let us mention that, since these theories are chiral, there are potential gauge anomalies for the $U(1)$ factors. Actually, even for the simplest choice of equal number of D4 branes in each box, the field theory is seemingly anomalous. This issue has been studied in [15], – in a T-dual picture of D1 branes at four-fold singularities, see section 5 –, where the anomaly was seen to be cancelled through an interaction with bulk modes. In the brane box picture, we expect some kind of anomaly inflow mechanism playing an analogous role. We leave this very interesting point for future research, and in the following will assume that such mechanism is at work and renders the theory consistent.

3.4. Examples

Field	**Representation**	**Field**
$H_{1,1,1}$	$(\square, \overline{\square}, 1, 1, 1, 1, 1, 1)$	$\overline{H}_{2,1,1}$
$H_{1,2,1}$	$(1, 1, \square, \overline{\square}, 1, 1, 1, 1)$	$\overline{H}_{2,2,1}$
$H_{1,1,2}$	$(1, 1, 1, 1, \square, \overline{\square}, 1, 1)$	$\overline{H}_{2,1,2}$
$H_{1,2,2}$	$(1, 1, 1, 1, 1, 1, \square, \overline{\square})$	$\overline{H}_{2,2,2}$
$V_{1,1,1}$	$(\square, 1, \overline{\square}, 1, 1, 1, 1, 1)$	$\overline{V}_{1,2,1}$
$V_{2,1,1}$	$(1, \square, 1, \overline{\square}, 1, 1, 1, 1)$	$\overline{V}_{2,2,1}$
$V_{1,1,2}$	$(1, 1, 1, 1, \square, 1, \overline{\square}, 1)$	$\overline{V}_{1,2,2}$
$V_{2,1,2}$	$(1, 1, 1, 1, 1, \square, 1, \overline{\square})$	$\overline{V}_{2,2,2}$
$N_{1,1,1}$	$(\square, 1, 1, 1, \overline{\square}, 1, 1, 1)$	$\overline{N}_{1,1,2}$
$N_{2,1,1}$	$(1, \square, 1, 1, 1, \overline{\square}, 1, 1)$	$\overline{N}_{2,1,2}$
$N_{1,2,1}$	$(1, 1, \square, 1, 1, 1, \overline{\square}, 1)$	$\overline{N}_{1,2,2}$
$N_{2,2,1}$	$(1, 1, 1, \square, 1, 1, 1, \overline{\square})$	$\overline{N}_{2,2,2}$
$D_{1,1,1}$	$(\square, 1, 1, 1, 1, 1, 1, \overline{\square})$	$\overline{D}_{2,2,2}$
$D_{2,1,1}$	$(1, \square, 1, 1, 1, 1, \overline{\square}, 1)$	$\overline{D}_{1,2,2}$
$D_{1,2,1}$	$(1, 1, \square, 1, 1, \overline{\square}, 1, 1)$	$\overline{D}_{2,1,2}$
$D_{2,2,1}$	$(1, 1, 1, \square, \overline{\square}, 1, 1, 1)$	$\overline{D}_{1,1,2}$

Table 3: Chiral multiplets for the $2\times2\times2$ box model. The ordering of the $U(n)$ factors in the second column is $U(n)_{1,1,1}\times U(n)_{2,1,1}\times U(n)_{1,2,1}\times U(n)_{2,2,1}\times U(n)_{1,1,2}\times U(n)_{2,1,2}\times U(n)_{1,2,2}\times U(n)_{2,2,2}$.

To see how the above set of rules works we consider an specific example with eight different three-dimensional boxes. Here we have two NS branes, two NS' branes and two NS" branes and an equal number n of D4 branes in each box. The theory is defined on $\mathbf{T}^3$ which arises from the grid identified by shifts by two boxes in each of the directions 246. Thus the $2\times2\times2$ unit cell consist of eight boxes, so the gauge group is $U(n)^8$. Even though the theory is $(0,2)$, the matter content is vector-like. The chiral multiplets in the model are listed in Table 3 (to make it shorter we list half or the fields and the conjugates of the other half)

In order to list the Fermi supermultiplets, we must make a choice of 'special' chiral superfield. For concreteness we will pick the fields D as the special ones. The corresponding Fermi multiplets are given in Table 4.

The interactions can be obtained by straightforward application of Eqs. (3.4) and (3.5).

Field	Representation	Field	Representation
$\Lambda^{(\overline{1})}_{1,1,1}$	$(\square, 1, 1, 1, 1, \overline{\square}, 1, 1)$	$\Lambda^{(\overline{1})}_{2,1,2}$	Conj. rep.
$\Lambda^{(\overline{1})}_{2,1,1}$	$(1, \square, 1, 1, \overline{\square}, 1, 1, 1)$	$\Lambda^{(\overline{1})}_{1,1,2}$	Conj. rep.
$\Lambda^{(\overline{1})}_{1,2,1}$	$(1, 1, \square, 1, 1, 1, 1, \overline{\square})$	$\Lambda^{(\overline{1})}_{2,2,2}$	Conj. rep.
$\Lambda^{(\overline{1})}_{2,2,1}$	$(1, 1, 1, \square, 1, 1, \overline{\square}, 1)$	$\Lambda^{(\overline{1})}_{1,2,2}$	Conj. rep.
$\Lambda^{(\overline{2})}_{1,1,1}$	$(\square, 1, 1, 1, 1, 1, \overline{\square}, 1)$	$\Lambda^{(\overline{2})}_{1,2,2}$	Conj. rep.
$\Lambda^{(\overline{2})}_{1,2,1}$	$(1, 1, \square, 1, \overline{\square}, 1, 1, 1)$	$\Lambda^{(\overline{2})}_{1,1,2}$	Conj. rep.
$\Lambda^{(\overline{2})}_{2,1,1}$	$(1, \square, 1, 1, 1, 1, 1, \overline{\square})$	$\Lambda^{(\overline{2})}_{2,2,2}$	Conj. rep.
$\Lambda^{(\overline{2})}_{2,2,1}$	$(1, 1, 1, \square, 1, \overline{\square}, 1, 1)$	$\Lambda^{(\overline{2})}_{2,1,2}$	Conj. rep.
$\Lambda^{(3)}_{1,1,1}$	$(\square, 1, 1, \overline{\square}, 1, 1, 1, 1)$	$\Lambda^{(3)}_{2,2,1}$	Conj. rep.
$\Lambda^{(3)}_{2,1,1}$	$(1, \square, \overline{\square}, 1, 1, 1, 1, 1)$	$\Lambda^{(3)}_{1,2,1}$	Conj. rep.
$\Lambda^{(3)}_{1,1,2}$	$(1, 1, 1, 1, \square, 1, 1, \overline{\square})$	$\Lambda^{(3)}_{2,2,2}$	Conj. rep.
$\Lambda^{(3)}_{2,1,2}$	$(1, 1, 1, 1, 1, \square, \overline{\square}, 1)$	$\Lambda^{(3)}_{1,2,2}$	Conj. rep.

Table 4: Fermi multiplets for the $2 \times 2 \times 2$ box model. The fields D have been chosen as the special chiral multiplets, and the Fermi superfields are of type $\Lambda^{(\overline{1})}$, $\Lambda^{(\overline{2})}$ and $\Lambda^{(3)}$. Recall that the ordering of the groups in the second column is $U(n)_{1,1,1} \times U(n)_{2,1,1} \times U(n)_{1,2,1} \times U(n)_{2,2,1} \times U(n)_{1,1,2} \times U(n)_{2,1,2} \times U(n)_{1,2,2} \times U(n)_{2,2,2}$.

4. Models with Enhanced Supersymmetry

4.1. Non-chiral enhancement of supersymmetry

In this section we discuss how to construct two-dimensional gauge theories with enhanced supersymmetry using the brane boxes introduced above. We begin by briefly mentioning the simpler case of non-chiral supersymmetry. We have already considered such theories in section 3.2, where we studied the dimensional reduction of four-dimensional $\mathcal{N} = 1$ models, *i.e.* $(2, 2)$ theories in two dimensions. It is easy to see that the pattern of the spectrum of the $(2, 2)$ field theories that can be constructed in our setup is that of the dimensional reductions of four-dimensional box models. Here by 'pattern' we mean the structure of the bi-fundamental multiplets, the quiver of the theory, regardless of the actual rank of the gauge factors. We stress this subtlety because, since $(2, 2)$ theories are non-chiral, there is more freedom in choosing the numbers of D4 branes in the boxes. So some of our $(2, 2)$ models would be dimensional reduction of anomalous four-dimensional theories. The point here has already been stressed in [21].

This relation to four-dimensional models greatly facilitates the construction of brane models with enhanced $(2,2)$, $(4,4)$ or $(8,8)$ supersymmetry. One basically takes the brane construction of four-dimensional $\mathcal{N}=1$, $\mathcal{N}=2$ or $\mathcal{N}=4$ theories in [6,10], replaces the D5 branes by D4 branes, and adds one NS$''$ brane. Thus for example, $(4,4)$ theories can be obtained by considering $k'=k''=1$, and arbitrary k. The unit cell is a finite row of k boxes, with trivial identifications of its sides. One such brane model is shown in Figure 7, for the particular case of $k=4$. The identification of the spectrum of the theory and the interactions can be done directly in four-dimensional language, using the rules in [6].

Similarly, gauge theories with sixteen supercharges can be obtained by considering a unit cell with a single box, with all sides identified. All fields in the theory are in the adjoint representation, and the brane configuration can be interpreted as D4 branes wrapped on a three-torus.



Fig. 7. The figure shows the brane box configuration which lead to a gauge theory in two dimensions with (4,4) supersymmetry. The unit cell consist of 4 boxes extending horizontally. Observe it has trivial identifications of its faces in all three directions 642. Recall that the numbers in the boxes are merely labels to distinguish them.

4.2. Chiral Supersymmetry Enhancement

(0,4) Theories

It is easy to construct brane box models with enhanced $(0,4)$ supersymmetry. The rule is that any model in which one kind of Fermi multiplet appears in the adjoint has at least $(0,4)$ supersymmetry. Instead of discussing it in general, let us present one such

example which illustrates the general features of these theories. The relevant facts about the field theory will be mentioned as needed (the essentials about $(0,4)$ supersymmetric theories can be extracted from [27]).

The configuration is the $4 \times 1 \times 1$ box model depicted in Figure 8. We label the boxes in the unit cell with *one* index i, and denote by n_i the number of D4 branes on each box. Notice that the faces of the unit cell are identified up to shifts in several directions. These non-trivial identifications of the sides of the unit cell can be read from the picture.



Fig. 8. The figure corresponds to a brane box configuration which leads to a two-dimensional field theory with (0,4) supersymmetry. The unit cell, also displayed, consists of 4 boxes extending along 6. Observe that in defining the three-torus by identifying the faces of the unit cell, the identifications are accompanied by shifts. These can be read from the figure. The model is said to have non-trivial identifications.

The gauge group is $\prod_i U(n_i)$. From our general rules we can read the matter content, which we have collected in Table 5 (the Fermi multiplets shown correspond to the choice of D as special field). We have also listed the $(0,4)$ multiplets these fields form.

For instance, the fields $\Lambda^{(3)}$, which transform in the adjoint, become part of the $(0,4)$ vector multiplet. These fermions, along with the $(0,2)$ gauginos, can be arranged in a four-dimensional vector, acted upon by the $SO(4)_R$ R-symmetry group of the field theory. Notice that only the usual $U(1)_R$ subgroup is visible in the brane configuration.

The fields H_i and V_{i+1} combine to form one $(0,4)$ chiral multiplet (since they transform in conjugate representations). The fields N_i and D_{i+2} form another $(0,4)$ chiral multiplet. The Fermi multiplets $\Lambda^{(\overline{1})}$ and $\Lambda^{(\overline{2})}$ remain as two $(0,4)$ Fermi multiplets, singlets under the R-symmetry.

$(\mathbf{0},\mathbf{2})$ **Field**	**Representation**	$(\mathbf{0},\mathbf{4})$ **Field**
H_i	$(\Box_i, \overline{\Box}_{i+1})$	Chiral
V_{i+1}	$(\Box_{i+1}, \overline{\Box}_i)$	
N_i	$(\Box_i, \overline{\Box}_{i+2})$	Chiral
D_{i+2}	$(\Box_{i+2}, \Box_i)$	
$\Lambda_i^{(\overline{1})}$	$(\Box_i, \overline{\Box}_{i+1})$	Fermi
$\Lambda_i^{(\overline{2})}$	$(\Box_i, \overline{\Box}_{i-1})$	Fermi
$\Lambda_i^{(3)}$	$(\Box_i, \overline{\Box}_i)$	Vector

Table 5: The table summarizes the matter content corresponding to the brane box configuration of Figure 8. The only field in the adjoint representation corresponds to a Fermi field. The rest of fields are charged as bifundamentals. The model has (0,4) supersymmetry, and in the last column we show how the $(0,2)$ fields combine in $(0,4)$ multiplets.

Concerning the interactions, they also respect $(0,4)$ supersymmetry. For instance, for the Fermi multiplet in the adjoint $\Lambda_i^{(3)}$ we have

$$\begin{aligned} J_i^{(3)} &= H_i V_{i+1} - V_i H_{i-1} \\ E_i^{(3)} &= D_i N_{i-2} - N_i D_{i+2}. \end{aligned} \tag{4.1}$$

It is easy to see that, when expressed in components, the Yukawa couplings from these terms are the 'gauge' interactions of the $(0,4)$ chiral multiplet with the additional gaugino $\lambda_i^{(3)}$ in the $(0,4)$ vector multiplet.

Let us also discuss the interactions of the remaining Fermi multiplets. The main novelty is that the two kinds of couplings of a Fermi multiplet to the chiral multiplets, given by the functions E and J, become related by supersymmetry. For instance, for the multiplet $\Lambda_i^{(\overline{1})}$, we have

$$\begin{aligned} J_i^{(\overline{1})} &= N_{i+1} H_{i+3} - H_{i+1} N_{i+2} \\ E_i^{(\overline{1})} &= D_i V_{i-2} - V_i D_{i-1}. \end{aligned} \tag{4.2}$$

The corresponding terms are the interaction of the Fermi multiplet $\Lambda^{(\overline{1})}$ with the $(0,4)$ multiplets formed by the pairs (H_i, V_{i+1}), and (N_i, D_{i+2}).

Analogously, the functions associated to the Fermi multiplet $\Lambda_i^{(\overline{2})}$,

$$\begin{aligned} J_i^{(\overline{2})} &= V_{i-1} N_{i-2} - N_{i-1} V_{i+1} \\ E_i^{(\overline{2})} &= D_i H_{i-2} - H_i D_{i+1}, \end{aligned} \tag{4.3}$$

contain the interactions of $\Lambda^{(2)}$ with the $(0,4)$ chiral multiplets.

This pattern holds in general. Any time there is one kind of Fermi multiplet in adjoint representations, the whole spectrum fits nicely into $(0,4)$ multiplets. It can also be checked that the interactions respect this supersymmetry. In order to avoid non-chiral supersymmetry enhancement there should not be any chiral multiplet in the adjoint, otherwise the supersymmetry would be $(4,4)$. There are many other models that can be constructed in a similar way, but we will not attempt a general classification. The supersymmetry enhancement of these models will receive a nice geometrical interpretation in section 5.4.

(0,6) Theories

It is also possible to get two-dimensional theories with six chiral supercharges. The appropriate way of obtaining them is constructing brane configurations which have two Fermi multiplets transforming in the adjoint. If the supersymmetry enhancement is to be chiral, one must also make sure that there are no chiral multiplets in adjoint representations.

In the following we present an example of one such model. The configuration is a $4 \times 1 \times 1$ box model, which we show in Figure 9. We label the boxes in the unit cell by an index i, as before.

The gauge group is $\prod_i U(n_i)$. The matter content is summarized in Table 6, where we also show how the $(0,2)$ fields combine to form $(0,6)$ multiplets.

The R-symmetry group is $SO(6)_R$. All four $(0,2)$ chiral multiplets combine into a single $(0,6)$ chiral multiplet. The four complex fields transform in the fundamental representation of $SU(4)_R \approx SO(6)_R$.

Also, $\Lambda^{(\overline{2})}$ and $\Lambda^{(3)}$, which transform in the adjoint, become part of the vector multiplet. Along with the gauginos in the $(0,2)$ multiplet, they can be arranged in a six dimensional real vector, acted upon by $SO(6)_R$

The field $\Lambda^{(\overline{1})}$ remains a Fermi multiplet. Again, the couplings also respect the higher supersymmetry. For instance the interactions of $\Lambda^{(\overline{2})}$ and $\Lambda^{(3)}$ become have the appropriate structure to be 'gauge' interactions.

It is easy to construct the most general brane box model with this structure. It is given by a generalization of the model in Figure 9, by considering a unit cell with an arbitrary number k of boxes. The spectrum is also given by Table 6 by simply increasing the range of variation of i, $i = 1, \ldots, k$.

$(\mathbf{0,2})$ **Field**	**Representation**	$(\mathbf{0,6})$ **Field**
H_i	$(\square_i, \overline{\square}_{i+1})$	Chiral
V_{i+1}	$(\square_{i+1}, \overline{\square}_i)$	
N_i	$(\square_i, \overline{\square}_{i+1})$	
D_{i+1}	$(\square_{i+1}, \overline{\square}_i)$	
$\Lambda_i^{(1)}$	$(\square_i, \overline{\square}_{i-2})$	Fermi
$\Lambda_i^{(2)}$	$(\square_i, \overline{\square}_i)$	Vector
$\Lambda_i^{(3)}$	$(\square_i, \overline{\square}_i)$	

Table 6: The table gives the matter content corresponding to the brane box configuration of Fig. 9. There are two Fermi multiplets in the adjoint representation. The rest of fields are charged as bifundamentals. It corresponds to a model with (0,8) supersymmetry.



Fig. 9. The figure shows a $4 \times 1 \times 1$ box model which leads to the construction of two-dimensional gauge theories with (0,6) supersymmetry. The unit cell consists of 4 boxes, and its faces are identified in a non-trivial way.

(0,8) Model

Finally, let us construct the only $(0,8)$ model which can be realized in our setup. The brane box configuration is given in Figure 10. The unit cell contains two boxes, and has non-trivial identifications of its faces. The gauge group is $U(n)^2$, and we easily check that all $(0,2)$ Fermi multiplets are in the adjoint representation. They become part of the $(0,8)$ gauge multiplet. The R-symmetry group $SO(8)_R$ acts on the eight-dimensional real vector formed by these fermions and the $(0,2)$ gauginos. Also, all $(0,2)$ chiral multiplets transform in the bifundamental $(\square_1, \overline{\square}_2)$ or its conjugate. The chiral multiplets will fill

$(0,8)$ chiral multiplets. In this case, all the $(0,2)$ interactions are 'gauge' interactions from the $(0,8)$ point of view. The basic features of $(0,8)$ theories have been determined *e.g.* in [29].



Fig. 10. The figure shows a $2 \times 1 \times 1$ box model giving a $(0,8)$ field theory. The unit cell contains two boxes, and its faces are identified in a non-trivial way.

The basic lesson we can learn from these further examples is that enhancement of supersymmetry is easily obtained in our setup. Non-chiral supersymmetries arise when one or several chiral multiplets transform in the adjoint of the gauge group. Enhancement of chiral supersymmetries appears when Fermi multiplets transform in the adjoint (and chiral multiplets do not). Notice that the chiral enhancement of supersymmetry is not manifest from the brane box point of view, and has to be checked by direct computation of the spectrum and interactions. In section 5.4 we will discuss how it becomes manifest in a T-dual configuration, where the field theory is realized in the world-volume of D1 branes at four-fold singularities. To establish such T-duality is the purpose of the following section.

5. The Interpretation of the Linear Sigma Model

In the previous sections we have introduced a large family of two-dimensional $(0,2)$ gauge theories. Since $(2,2)$ and $(0,2)$ theories have been traditionally used as world-sheet descriptions of string theories propagating on some target space, it is a natural question whether the (classical) Higgs branch or our models has any geometrical interpretation of

the kind. In this section we are to show that it describes the dynamics of a type IIB D1 brane on a $\mathbf{C}^4/\Gamma$ singularity, with Γ an abelian subgroup of $SU(4)$. The main tool for reaching this conclusion will be a T-duality performed on the brane box model along the directions 246.

5.1. T-duality Along 246

In this section we perform a T-duality on the brane box models along the directions 246. The main tool will be the well known T-duality relation between a set of n parallel NS fivebranes and n Kaluza-Klein monopoles. The discussion in this subsection parallels that in [12].

Let us start with the simplest case of a brane box model formed by a unit cell of $k \times k' \times k''$ boxes, with trivial identifications of faces. In this case the T-duality along the directions 246 is particularly easy. We start with k NS branes along 012345, k' NS$'$ branes along 012367, and k'' NS$''$ branes along 014567. The T-duality along 2 transforms the NS$''$ branes into k'' Kaluza-Klein monopoles. These will be described by a multi-center Taub-NUT metric, with non-trivial geometry on the directions 2$'$,3,8,9, with 2$'$ denoting the coordinate dual to 2. Notice that, since the 3,8,9 coordinates of the initial NS$''$ branes coincided, so do the coordinates of the corresponding k'' centers of the Taub-NUT metric, so that it contains singularities of type $A_{k''-1}$.

Similarly, the T-duality along 4 transforms the k' NS$'$ branes into k' Kaluza-Klein monopoles with world-volume along 012367, and represented by a nontrivial geometry on 4$'$,5,8,9. Again, since the centers of the Kaluza-Klein monopoles coincide, such geometry will contain singularities of type $A_{k'-1}$. Finally, the T-duality along 6 turns the k NS branes into k Kaluza-Klein monopoles. Their world-volume spans 012345, and they are represented by a non-trivial geometry along 6$'$,7,8,9. Since again all the centers coincide, there will be A_{k-1} singularities.

Thus, the final T-dual of the grid of NS, NS$'$ and NS$''$ branes is type IIB string theory with a complicated geometry in the directions 2$'$,3,4$'$,5,6$'$,7,8,9. One can think of it roughly as some 'superposition' of the Kaluza-Klein monopoles we have described. Even without a quantitative knowledge of such metric, we can describe the relevant features for our purposes. One such feature is that the number of unbroken supersymmetries constrains the manifold to be a Calabi-Yau four-fold. Also, from our remarks above we know the existence of certain (complex) surfaces of singularities of type A_{k-1}, $A_{k'-1}$ and $A_{k''-1}$

singularities. If we introduce complex coordinates $w_1 = \exp(x^7 + ix^{6'})$,$w_2 = \exp(x^5 + ix^{4'})$, $w_3 = \exp(x^3 + ix^{2'})$, and $w_4 = x^9 + ix^8$, the surface of A_{k-1} singularities is defined roughly by $w_1 = w_4 = 0$, the surface of $A_{k'-1}$ singularities is defined by $w_2 = w_4 = 0$, and the surface of $A_{k''-1}$ singularities is given by $w_3 = w_4 = 0$. At the origin $w_1 = w_2 = w_3 = w_4 = 0$ all surfaces meet and the singularity is worse. It can be described as a quotient singularity of type $\mathbb{C}^4/\Gamma$, with $\Gamma = \mathbb{Z}_k \times \mathbb{Z}_{k'} \times \mathbb{Z}_{k''}$. This discrete group is generated by elements θ, ω, η, whose action on $(z_1, z_2, z_3, z_4) \in \mathbb{C}^4$ is as follows:

$$\begin{array}{rccc} \theta: & (z_1, z_2, z_3, z_4) & \rightarrow & (e^{2\pi i/k} z_1, z_2, z_3, e^{-2\pi i/k} z_4) \\ \omega: & (z_1, z_2, z_3, z_4) & \rightarrow & (z_1, e^{2\pi i/k'} z_2, z_3, e^{-2\pi i/k'} z_4) \\ \eta: & (z_1, z_2, z_3, z_4) & \rightarrow & (z_1, z_2, e^{2\pi i/k''} z_3, e^{-2\pi i/k''} z_4). \end{array} \tag{5.1}$$

In this description it becomes clear that there may be further surfaces of singularities when the greatest common divisor of any two of k, k', k'' is not 1, in analogy with the discussion in [12]. This will not be relevant for our purposes and we do not develop the issue further.

After the T-duality, the initial D4 branes become D1 branes located at a point in the four-fold. When the initial D4 branes are bounded by the grid of NS, NS$'$ and NS$''$ branes, the T-dual D1 branes will be located precisely at the $\mathbb{C}^4/\Gamma$ singular point. The field theories introduced in the previous sections correspond to the field theories appearing in the world-volume of such D1 brane probes. In the following subsection we will show how the structure of the singularity controls the spectrum and dynamics of the field theory.

5.2. The spectrum of the brane at the singularity

The field theory on the world-volume of a D brane probes on singularities can be determined using open string techniques introduced in [13,16]. The particular case of D1 branes on four-fold quotient singularities $\mathbb{C}^4/\Gamma$ has been studied in [15], whose results we review in the present section. We will center on *abelian* discrete groups, and we will also show how the spectra obtained in this case actually match with those we presented in Section 3. This will confirm the spectra we had proposed, and also illustrate the usefulness of the T-duality between brane boxes and branes at singularities.

Let us briefly review the field theory on the world-volume of N D1 branes on $\mathbb{C}^4$. This is the dimensional reduction of ten-dimensional $\mathcal{N} = 1$ $U(N)$ super Yang-Mills to two dimensions. The R-symmetry group is $SO(8)_R$, which can be understood as the symmetry group of the transversal dimensions. The theory contains a set of gauge bosons A_μ, singlets

under the R-symmetry. The matter fields are eight real scalar fields ϕ transforming in the $\mathbf{8_v}$ of $SO(8)$, eight left-handed fermions ψ_- transforming in the $\mathbf{8_s}$, and eight right-handed fermions ψ_+, which transform in the $\mathbf{8_c}$. All fields are in the adjoint representation of the gauge group.

The field theory of D1 branes on $\mathbb{C}^4/\Gamma$ can be obtained as a projection of the theory just described onto Γ invariant states. The group Γ acts on the R-symmetry of the theory, since this symmetry associated to the transversal coordinates, which are modded out to form the quotient. Since Γ is a subgroup of $SU(4)$, an unbroken $U(1)_R$ R-symmetry will remain in the quotient theory, due to the decomposition $SO(8)_R \supset SU(4) \times U(1)_R$. This action of Γ on $\mathbb{C}^4$ is specified by a four-dimensional faithful representation, which we will denote by $\mathbf{4}$. Since all irreducible representations of Γ are one-dimensional (recall we only consider the abelian case), the representation $\mathbf{4}$ is reducible, and has a decomposition

$$\mathbf{4} = \mathcal{R}_{A_1} \oplus \mathcal{R}_{A_2} \oplus \mathcal{R}_{A_3} \oplus \mathcal{R}_{A_4} \tag{5.2}$$

with the restriction that the tensor product of all representations is the trivial representation. This is required so that $\mathbf{4}$ defines an action in $SU(4)$, rather than in $U(4)$. It also can be expressed as $A_4 = -A_1 - A_2 - A_3$, in notation of [12].

For future convenience, let us point out the this choice of $\mathbf{4}$ also determines a six-dimensional representation $\mathbf{6}$, which is obtained by taking the antisymmetric part of $\mathbf{4} \otimes \mathbf{4}$

$$\mathbf{6} = \mathcal{R}_{A_1 \oplus A_2} \oplus \mathcal{R}_{A_1 \oplus A_3} \oplus \mathcal{R}_{A_2 \oplus A_3} \oplus \mathcal{R}_{-A_1 - A_2} \oplus \mathcal{R}_{-A_1 - A_3} \oplus \mathcal{R}_{-A_2 - A_3} \tag{5.3}$$

where we have introduced the notation $\mathcal{R}_{A_i} \otimes \mathcal{R}_{A_j} = \mathcal{R}_{A_i \oplus A_j}$ for the tensor product of representations. The action of Γ must also be embedded in the Chan-Paton factors of the D1 branes. This is specified by giving a representation of Γ which we will denote by $\mathcal{R}_{C.P.}$. The configuration of the D1 branes is essentially described by such representation.

The simplest physical situation is that of n D1 branes in the quotient manifold. This can be equivalently described in the cover $\mathbb{C}^4$ of the orbifold space as a set of $n|\Gamma|$ D1-branes (with $|\Gamma|$ being the order of the discrete group) with Chan-Paton factors in n copies of the adjoint representation $\mathcal{R}_\Gamma$,

$$\mathcal{R}_{C.P.} = n\mathcal{R}_\Gamma = n \oplus_I \mathcal{R}_I. \tag{5.4}$$

The fields in the resulting $(0,2)$ field theory on the D1 branes will be those invariant under the simultaneous action on R-symmetry and gauge quantum numbers. Following

the rules in [13] we can obtain the spectrum. The gauge group is $U(n)^{|\Gamma|} = \prod_I U(n)_I$, so there is one gauge factor associated to each irreducible representation of Γ.

Since the scalar fields transform in the $\mathbf{8_v}$ of $SO(8)_R$, which decomposes as $\mathbf{4}+\overline{\mathbf{4}}$ of the SU(4), to compute the scalar spectrum in the orbifold theory we have to compute the tensor products of the representation $\mathbf{4}$ (5.2) with each irreducible representation $\mathcal{R}_I$,

$$\mathbf{4} \otimes \mathcal{R}_I = \mathcal{R}_{I\oplus A_1} \oplus \mathcal{R}_{I\oplus A_2} \oplus \mathcal{R}_{I\oplus A_3} \oplus \mathcal{R}_{I-A_1-A_2-A_3} \tag{5.5}$$

This implies that there are four kinds of complex scalar fields. The first, associated to the first complex plane in $\mathbb{C}^4$, transform as the $(\square, \overline{\square})$ of $U(n)_I \times U(n)_{I\oplus A_1}$; we denote these fields as $\phi_{I,I\oplus A_1}$. The second corresponds to fields denoted $\phi_{I,I\oplus A_2}$, transforming as the $(\square, \overline{\square})$ of $U(n)_I \times U(n)_{I\oplus A_2}$. There are fields $\phi_{I,I\oplus A_3}$, which transform in the $(\square, \overline{\square})$ of $U(n)_I \times U(n)_{I\oplus A_3}$. Finally, there are fields $\phi_{I,I-A_1-A_2-A_3}$ transforming in the $(\square, \overline{\square})$ of $U(n)_I \times U(n)_{I-A_1-A_2-A_3}$.

The right-handed fermions ψ_+ transform in the $\mathbf{8_c}$ of $SO(8)_R$, which also decomposes as $\mathbf{4}+\overline{\mathbf{4}}$ under $SU(4)$. To obtain the invariant fields in the orbifold theory one has to use again the products of the $\mathbf{4}$ with the representations $\mathcal{R}_I$, (5.5). Thus one obtains four kinds of such fermions fields, denoted $\psi_{I,I\oplus A_i}$, for $i = 1, \ldots, 4$, which transform in the $(\square, \overline{\square})$ of $U(n)_I \times U(n)_{I\oplus A_i}$. The scalars $\phi_{I,I\oplus A_i}$ and the right-handed fermions $\psi_{I,I\oplus A_i}$ form together $(0,2)$ chiral multiplets, denoted $\Phi^i{}_I$ in the following.

The left-handed fermions ψ_- transform in the $\mathbf{8_s}$ of $SO(8)_R$, whose decomposition under $SU(4)$ is $\mathbf{8_s} = \mathbf{6}+\mathbf{1}+\mathbf{1}$. The two fermions transforming trivially under the $SU(4)$ will give rise to one left-handed fermion χ_I (and its conjugate) transforming in the adjoint of the gauge factor $U(n)_I$. These fermions, together with the gauge bosons, form $(0,2)$ vector multiplets. On the other hand, we have to compute the products of the representation $\mathbf{6}$ (5.3) with the representations $\mathcal{R}_I$, in order to get the remaining left-handed fermions. This yields

$$\begin{aligned} \mathbf{6} \otimes \mathcal{R}_I = & \mathcal{R}_{I\oplus A_1\oplus A_2} \oplus \mathcal{R}_{I\oplus A_1\oplus A_3} \oplus \mathcal{R}_{I\oplus A_2\oplus A_3} \\ & \oplus \mathcal{R}_{I-A_1-A_2} \oplus \mathcal{R}_{I-A_1-A_3} \oplus \mathcal{R}_{I-A_2-A_3}. \end{aligned} \tag{5.6}$$

Thus we obtain left-handed fermions $\psi_{I,I\oplus A_i\oplus A_j}$, for $a, b = 1, \ldots, 4$, $i \neq j$, transforming in the $(\square, \overline{\square})$ of $U(n)_I \times U(n)_{I\oplus A_i\oplus A_j}$. These will correspond to three Fermi multiplets and their conjugates. We will denote them by Λ^{ij}_I. Again, which fields are considered the Fermi multiplets, and which their conjugates, depends on the choice of a 'special' chiral

multiplet, to appear in the functions E of the Fermi multiplets. Following the convention in [15], we take Φ^4_I as these special fields. This implies that the fields Λ^{i4}_I are the Fermi multiplets, and that Λ^{ij}_I, $i,j \neq 4$ are the conjugates.

The interactions are also given by the projection of the interactions in the non-orbifolded theory onto Γ invariant terms. The outcome can be recast as a definition of functions E and J for the Fermi multiplets. The functions E are

$$\begin{array}{lcll} \Lambda^{14}_I & \to & E^1_I & = \Phi^4_I \Phi^1_{I-A_1-A_2-A_3} - \Phi^1_I \Phi^4_{I+A_1} \\ \Lambda^{24}_I & \to & E^2_I & = \Phi^4_I \Phi^2_{I-A_1-A_2-A_3} - \Phi^2_I \Phi^4_{I+A_2} \\ \Lambda^{34}_I & \to & E^3_I & = \Phi^4_I \Phi^3_{I-A_1-A_2-A_3} - \Phi^3_I \Phi^4_{I+A_3}. \end{array} \tag{5.7}$$

And the functions J are

$$\begin{array}{lcll} \Lambda^{14}_I & \to & J^1_I & = \Phi^2_{I-A_2-A_3} \Phi^3_{I-A_3} - \Phi^3_{I-A_2-A_3} \Phi^2_{I-A_2} \\ \Lambda^{24}_I & \to & J^2_I & = \Phi^3_{I-A_1-A_3} \Phi^1_{I-A_1} - \Phi^1_{I-A_1-A_3} \Phi^3_{I-A_3} \\ \Lambda^{34}_I & \to & J^3_I & = \Phi^1_{I-A_1-A_2} \Phi^2_{I-A_2} - \Phi^2_{I-A_1-A_2} \Phi^1_{I-A_1}. \end{array} \tag{5.8}$$

Observe that the condition $\sum_{i,I} J^i_I E^i_I = 0$ is verified. The interaction terms in components are easily obtained using the equations in Section 2.

5.3. Comparison with the Brane Box Models

In this section we analyze the particular case of $\Gamma = \mathbb{Z}_k \times \mathbb{Z}_{k'} \times \mathbb{Z}_{k''}$, with the generators θ, ω, η of $\mathbb{Z}_k$, $\mathbb{Z}_{k'}$, $\mathbb{Z}_{k''}$ acting on $\mathbb{C}^4$ as in (5.1). In this case the group has $|\Gamma| = k \cdot k' \cdot k''$ irreducible representations, $\mathcal{R}_{a,b,c}$, with a, b, c integers defined modulo k, k', k'', respectively. The representation $\mathcal{R}_{a,b,c}$ associates to the group element $\theta^l \omega^m \eta^n$ the phase factor $\exp\left[\, 2\pi i (\frac{al}{k} + \frac{bm}{k'} + \frac{cn}{k''}) \,\right]$. Clearly, the tensor product of representations is given by $\mathcal{R}_{a,b,c} \otimes \mathcal{R}_{a',b',c'} = \mathcal{R}_{a+a',b+b',c+c'}$, *i.e.* separated addition in the indices.

With this definition, the action of Γ on $\mathbb{C}^4$ corresponds to the choice

$$\mathbf{4} = \mathcal{R}_{1,0,0} \oplus \mathcal{R}_{0,1,0} \oplus \mathcal{R}_{0,0,1} \oplus \mathcal{R}_{-1,-1,-1}. \tag{5.9}$$

From it we can also obtain

$$\mathbf{6} = \mathcal{R}_{1,1,0} \oplus \mathcal{R}_{1,0,1} \oplus \mathcal{R}_{0,1,1} \oplus \mathcal{R}_{-1,-1,0} \oplus \mathcal{R}_{-1,0,-1} \oplus \mathcal{R}_{0,-1,-1}. \tag{5.10}$$

Using the rules we have described above we can easily find out the spectrum of the theory. The gauge group is $U(n)^{kk'k''} = \prod_{a,b,c} U(n)_{a,b,c}$, where each factor is associated to an irreducible representation. Let us now look at the four types of $(0,2)$ chiral multiplets

$\Phi_{I,I\oplus A_i}$, $i=1,\ldots,4$. The fields associated to the first complex plane, $\Phi^1_{a,b,c}$, transform in the $(\square,\overline{\square})$ of $U(n)_{a,b,c}\times U(n)_{a+1,b,c}$. The fields associated to A_2, $\Phi^2_{a,b,c}$, transform in the $(\square,\overline{\square})$ of $U(n)_{a,b,c}\times U(n)_{a,b+1,c}$. The fields $\Phi^3_{a,b,c}$ transform in the $(\square,\overline{\square})$ of $U(n)_{a,b,c}\times U(n)_{a,b,c+1}$. Finally, the fields $\Phi^4_{a,b,c}$ transform in the $(\square,\overline{\square})$ of $U(n)_{a,b,c}\times U(n)_{a-1,b-1,c-1}$.

The Fermi multiplets are obtained using the representation **6** above. The field $\Lambda^{14}_{a,b,c}$ transforms in the $(\square,\overline{\square})$ of $U(n)_{a,b,c}\times U(n)_{a,b-1,c-1}$; the fields $\Lambda^{24}_{a,bc}$ transform in the $(\square,\overline{\square})$ of $U(n)_{a,b,c}\times U(n)_{a-1,b,c-1}$; the fields $\Lambda^{34}_{a,b,c}$ transform in the $(\square,\overline{\square})$ of $U(n)_{a,b,c}\times U(n)_{a-1,b-1,c}$.

We have listed the spectrum of chiral multiplets and Fermi multiplets in Table 7.

Field	**Representation**	**Brane box**
$\Phi^1_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a+1,b,c}$	$H_{a,b,c}$
$\Phi^2_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a,b+1,c})$	$V_{a,b,c}$
$\Phi^3_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a,b,c+1})$	$N_{a,b,c}$
$\Phi^4_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a-1,b-1,c-1})$	$D_{a,b,c}$
$\Lambda^{14}_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a,b-1,c-1})$	$\Lambda^{(\overline{2})}_{abc}$
$\Lambda^{24}_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a-1,b,c-1})$	$\Lambda^{(\overline{1})}_{a,b,c}$
$\Lambda^{34}_{a,b,c}$	$(\square_{a,b,c},\overline{\square}_{a-1,b-1,c})$	$\Lambda^{(3)}_{a,b,c}$

Table 7: The matter content of the theory of D1 branes at singularities. In the last column we compare these fields with the spectrum obtained in a $k\times k'\times k''$ brane box model of Section 3.

Notice that the spectrum we have obtained corresponds precisely to the one we proposed for a $k\times k'\times k''$ box model with trivial identifications of sides. The chiral multiplets $\Phi^i_{a,b,c}$ correspond to the fields H, V, N, D of Section 3. The choice of Φ^4 as special field corresponds to our convention of section 3.3, of choosing the D fields as special. The Fermi multiplets Λ^{14}, Λ^{24}, Λ^{34} in the singularity correspond to the Fermi fields $\Lambda^{(\overline{2})}$, $\Lambda^{(\overline{1})}$, $\Lambda^{(3)}$, respectively. This comparison is shown in the last column of Table 7. It is also a simple matter to particularize expressions (5.7) and (5.8) to this case and reproduce the interactions proposed in Section 3.3

The matching of the two field theories was actually expected, since we had already argued that the configurations are related by T-duality. The detailed correspondence of the spectra and interactions in both constructions is a nice argument supporting both the T-duality, and the spectrum proposed in Section 3 for the brane box configuration.

The T-duality proposal can be generalized in several directions. For instance, there is no difficulty in extending it to models with non-trivial identifications of faces of the unit cell, following [12]. Also, by considering fractional branes at the singularity one finds the T-duals of brane box models with different number of D4 branes in each box.

Rather than commenting on these exercises, we turn to briefly mention a few facts about the phase structure (in the sense of [24]) of these field theories as linear sigma models. Even though the field theories are relatively complicated, this analysis has been carried out for $(2,2)$ theories [16], and $(0,2)$ theories [15][7]. The outcome is that only the geometric phases of the linear sigma model are realized, and so no CY/LG correspondence holds in this case. However, flop transitions between different geometric phases [30] are certainly possible [31]. Since the field theories in this section are the same as those arising from brane box models (via the T-duality map), the results mentioned above also apply in the brane box context.

In the following Section we illustrate another interesting application of the T-duality map. We turn to the study of the models with enhanced supersymmetries introduced in Section 4.

5.4. Enhanced Supersymmetries

When the discrete group Γ that defines the singularity is not a generic subgroup of $SU(4)$, so that a larger R-symmetry is left unbroken, we can expect an enhancement of supersymmetry. The simplest such examples, yielding non-chiral supersymmetry enhancement appear when $\Gamma \subset SU(3)$. The R-symmetry of the theory is $U(1) \times U(1)$, and the theories obtained are $(2,2)$ in two dimensions. These models are clearly T-dual of configurations of D3 branes at Calabi-Yau three-fold singularities $\mathbb{C}^3/\Gamma$, which yield four-dimensional $\mathcal{N} = 1$ theories. Analogously, when $\Gamma \subset SU(2)$, the $(4,4)$ theories obtained are related of six-dimensional $\mathcal{N} = 1$ theories, which can be realized in the world-volume of D5 branes at ALE singularities $\mathbb{C}^2/\Gamma$.

The relation with the brane box model is clear. When the singularity has a restricted holonomy group, $SU(3)$ (resp. $SU(2)$), the brane box configuration reproducing the $(2,2)$ (resp. $(4,4)$) field theory has only two (resp. one) kinds of NS fivebranes. Thus the brane box configurations also admit an interpretation as reduction of higher dimensional brane models.

[7] The theories in these references actually correspond to the case where all the gauge factors are $U(1)$'s.

More interesting is the chiral enhancement of supersymmetry. As mentioned in [15], one can obtain $(0,4)$ theories in the D1 brane picture when the four-fold singularity admits some additional covariantly constant spinors. This can be translated in the condition $\Gamma \subset SU(2) \times SU(2)$, which leaves a $SU(2) \times SU(2)$ unbroken R-symmetry. This restricted holonomy implies the Γ-invariance of the forms $dz_1 \wedge dz_2$ and $dz_3 \wedge dz_4$ in $\mathbb{C}^4$. This condition restricts the possible actions of Γ on $\mathbb{C}^4$, in such a way that it ensures the existence of one Fermi multiplet in the adjoint. This fermions are required in a $(0,4)$ theory since they become part of the $(0,4)$ vector multiplet. Also, one can check that the chiral multiplets arrange in conjugate pairs to form $(0,4)$ chiral multiplets. The interactions also respect the higher supersymmetry.

These and some of the following special holonomies have also appeared in the study of two-dimensional compactifications of string theory and M-theory (see for example [28]). A particular family of such singularities is mentioned in [15]. The group is $\mathbb{Z}_n$, with the generator θ acting on $\mathbb{C}^4$ as

$$\theta : \ (z_1, z_2, z_3, z_4) \ \rightarrow \ (e^{2\pi i/n} z_1, e^{-2\pi i/n} z_2, e^{2\pi i a/n} z_3, e^{-2\pi i a/n} z_4) \tag{5.11}$$

It is a simple matter to obtain the spectrum and interactions in this example. One can also easily find brane box configurations yielding the same field theory. In fact, the example shown in Figure 8 correspond to the $\mathbb{Z}_4$ $n = 4, a = 2$ model in the family above.

Another interesting family of such theories is obtained as the product of two ALE singularities living in the (z_1, z_2) and (z_3, z_4) spaces, respectively.

Let us stress that for *any* singularity yielding a $(0,4)$ theory we can find brane box configurations reproducing the same field theory [8]. The construction is systematic and closely analogous to that discussed in [12] for three-fold singularities, so we will not discuss it here.

Observe that the basic requirement in order to have $(0,4)$ supersymmetry can be rephrased as the existence of Fermi multiplets in the adjoint representation. This agrees with our comments from the brane box point of view in section 4.2.

Concerning the field theories with $(0,6)$ supersymmetry, they are also obtained using singularities with restricted holonomy. In this case the condition amounts to ensuring the existence of two $(0,2)$ Fermi multiplets in the adjoint representation (and no adjoint chiral

[8] As in [12], there are in general several different brane box models corresponding to the same field theory. They correspond to T-dualizing the same singularity along different sets of directions.

multiplets). A complete classification of such singularities is possible, the only such models are $\mathbb{Z}_n$ orbifolds with the generator θ acting as

$$\theta : \ (z_1, z_2, z_3, z_4) \ \rightarrow \ (e^{2\pi i/n} z_1, e^{-2\pi i/n} z_2, e^{2\pi i/n} z_3, e^{-2\pi i/n} z_4). \tag{5.12}$$

The unbroken R-symmetry is $SU(4)_R \approx SO(6)_R$. Again it is a simple exercise to obtain the spectrum and interactions of the theory, and find brane box models yielding the same field theory. As an example, let us mention that the four-box model shown in Figure 9 gives precisely the field theory of the $\mathbb{Z}_4$ example in the family just mentioned. The general $\mathbb{Z}_n$ case is obtained by considering an analogous brane box model with n boxes. As indicated in section 4.2, the spectrum in the general case is given in Table 6.

Finally, let us comment on the only singularity yielding a $(0,8)$ supersymmetric theory. It is the $\mathbb{Z}_2$ orbifold whose generator θ acts as inversion of the four coordinates in $\mathbb{C}^4$. The unbroken R-symmetry is $SO(8)_R$. As for the spectrum, the gauge group is $U(n)^2$, all the Fermi multiplets transform in the adjoint, and all chiral multiplets transform in bi-fundamental representations. The model is clearly the same one we discussed in section 4.2, and corresponds to the brane box configuration shown in Figure 10.

The main conclusion of this discussion is that enhanced supersymmetry is associated to a restricted holonomy in the four-fold singularity. The T-duality map discussed in previous subsection then seems to suggest that the T-dual brane configurations present enhanced supersymmetry due to properties of their 'generalized holonomy group' in the sense of [20]. In the cases of non-chiral supersymmetry enhancement this is obvious, since essentially the correspond to removing one kind of brane. In the more interesting case of chiral supersymmetries, on the other hand, the supersymmetry enhancement is not so obvious in the brane box configuration and only follows upon the detailed computation of spectrum and interactions. It would be nice to achieve a better understanding of such brane box models.

6. Concluding Remarks

In this paper we have introduced brane box configurations giving chiral field theories in two dimensions. An interesting feature is that are a natural generalization of brane models

yielding chiral field theories in six and four dimensions. The connection between the chiral theories in the different dimensions is dimensional reduction followed by an appropriate chiral projection, performed by the introduction of a new kind of NS fivebrane.

We have provided the rules to determine the field theory spectrum from the brane box data. Since the theories have only two supersymmetries, the interactions have a relatively complicated structure. However, we have shown how they can be easily encoded in a diagrammatic representation in the brane box configuration.

We have also discussed the conditions under which the models present enhanced supersymmetry. We have stressed that it is possible to construct brane box models with enhanced $(0,4)$, $(0,6)$ and $(0,8)$ supersymmetry, and shown several examples of such configurations.

In the case where the direction 246 in the brane configuration are compact, we have related the corresponding models to theories arising from D1 brane probes at $\mathbb{C}^4/\Gamma$ singularities, with Γ an abelian subgroup of $SU(4)$. The relation is based on T-duality along the compact directions. This result has been useful, since it provides a simple rederivation of the rules to compute the spectrum and interactions we had proposed in the brane box picture. It also provides a nice geometrical interpretation for the enhancement of supersymmetry (both chiral and non-chiral) in terms of the holonomy group of the four-fold. Much could be said about the relation of the brane boxes and the singularity theories. However, since this discussion is essentially identical to that in [12], we have not repeated the details. Let us stress however that there is no difficulty in repeating the exercise in the four-fold case.

There are several interesting results concerning the phase structure of these field theories, when understood as linear sigma models [16,15]. There the field theories appeared from D-branes at singularities. An interesting application of the T-duality map is that it ensures these are the same theories one obtains from brane box configurations. This allows us to directly borrow the corresponding results and learn that only the geometric phases are accesible to the field theories. However, flop phase transition are still possible, and it would be interesting to give them a direct intepretation in the brane box picture.

Another interesting point is that this relation between brane boxes and branes at singularities is expected to shed light on some issues which are not completely clear in the brane box construction, such as the restrictions in the number of D4 branes that one is allowed to put in the boxes. Such restrictions are required to ensure cancellation of

gauge anomalies, and can be presumably studied in the singularity picture in the context of perturbative string theory. We leave this as an open question.

Observe that the singularity picture provides the construction of a larger family of $(0,2)$ theories, by considering non-abelian subgroups of $SU(4)$. The rules to compute the spectrum are a simple generalization of those we mentioned in the abelian case.

Finally let us mention an amusing relation between the $(0,2)$ theories we have studied and the dimensional reduction of the four-dimensional non-supersymmetric orbifold theories introduced in [19]. As we have mentioned, the theory of D1 branes on $\mathbb{R}^8$ has $SO(8)_R$ R-symmetry. The scalar fields ϕ transform in the $\mathbf{8_v}$ representation, the right-handed spinors ψ_+ transform in the $\mathbf{8_c}$, and the left-handed spinors transform in the $\mathbf{8_s}$.

There are three inequivalent embeddings of $SU(4)$ in $SO(8)$. They differ in which of the three representations '$\mathbf{8}$' decomposes as $\mathbf{6}+\mathbf{1}+\mathbf{1}$. Since the three representations are related by $SO(8)$ triality, so are the three embeddings. Taking this into account, given a (possibly non-abelian) discrete group $\Gamma \subset SU(4)$, and four- and six- dimensional representations $\mathbf{4}$ and $\mathbf{6}$, there are three possible orbifold theories we can define, using the three embeddings of $SU(4)$ in $SO(8)$.

We have already met one of these, where $\mathbf{8}_s$ decomposes as $\mathbf{6}+\mathbf{1}+\mathbf{1}$, and both $\mathbf{8_v}$ and $\mathbf{8_c}$ decompose as $\mathbf{4}+\overline{\mathbf{4}}$. The orbifold theory is a $(0,2)$ chiral supersymmetric theory. It contains scalars $\phi_{I,I\oplus A_i}$, right-handed fermions $\psi^+_{I,I\oplus A_i}$, and left-handed fermions $\psi^-_{I,I\oplus A_i\oplus A_j}$ (in notation of Section 5). The right-handed spinors and scalars are related by the right-handed supersymmetries. The conmutant of the $SU(4)$ is the $U(1)$ R-symmetry of the theory.

Another embedding, in which $\mathbf{8}_c$ decomposes as $\mathbf{6}+\mathbf{1}+\mathbf{1}$, and $\mathbf{8}_v$, $\mathbf{8}_s$ decompose as $\mathbf{4}+\overline{\mathbf{4}}$, is clearly the parity transformed of the precedent. The scalars $\phi_{I,I\oplus A_i}$ and the left-handed fermions $\psi^-_{I,I\oplus A_i}$ are related by $(2,0)$ supersymmetry. There are also right-handed fermions $\psi^+_{I,I\oplus A_i\oplus A_j}$.

Finally there is a third choice, in which $\mathbf{8}_v$ decomposes as $\mathbf{6}+\mathbf{1}+\mathbf{1}$, and $\mathbf{8}_s$, $\mathbf{8}_c$ decompose as $\mathbf{4}+\overline{\mathbf{4}}$. The field theory contains real scalars $\phi_{I,I\oplus A_i\oplus A_j}$, complex scalars $\phi_{I,I}$, and vector-like fermions $\psi^\pm_{I,I\oplus A_i}$. These theories are the dimensional reduction of the four-dimensional non-supersymmetric theories introduced in [19]. Now the $U(1)$ which is unbroken arises in the dimensional reduction. Observe the spectrum is similar to that of the $(0,2)$ or $(2,0)$ theories by an exchange of the scalars with one kind of fermions. This is somewhat reminiscent of the twisting of supersymmetric theories where one mixes the Lorentz group and the internal symmetries of the theory, even though we have not

been able to make this more precise. It is tantalizing to imagine that this amusing group theoretical fact may have some deeper meaning.

Acknowledgements

It is a pleasure to thank A. Hanany and R. Blumenhagen for useful comments. H. G-C. thanks to E. Witten for hospitality in the Institute for Advanced Study. A. U. is indebted to M. González for her support. The work H. G-C. is supported by a Postdoctoral CONACyT fellowship (Mexico) under the program *Programa de Posdoctorantes: Estancias Posdoctorales en el Extranjero para Graduados en Instituciones Nacionales 1996-1997.* The work of A. U. is supported by a fellowship from the Ramón Areces Foundation (Spain), and partially by the CICYT (Spain), under grant AEN97-1678.

Appendix I. Parameters in the Model: Gauge Couplings, Theta Parameters, Fayet-Iliopoulos Parameters

Gauge couplings, theta angles and Fayet-Iliopoulos (FI) parameters can be characterized in analogy with the four-dimensional brane box configurations [6,10].

Gauge couplings

The gauge couplings of the various gauge groups are given in terms of the positions of the NS branes along x_6 direction, the positions of the NS' branes along the x_4 direction and the positions of the NS" branes along the x^2 direction. There are k possible positions x_6^i, k' positions x_4^j and k'' x_2^k. These directions are divided into k intervals of lengths $I_a = x_6^a - x_6^{a-1}$ for x_6, into k' intervals of lengths $J_b = x_4^b - x_4^{b-1}$ for x_4 and into k'' intervals of lengths $K_c = x_2^c - x_2^{c-1}$ for x_2. The intervals I_a, J_b and K_c satisfy the relations $\sum_a I_a = R_6$, $\sum_b J_b = R_4$ and $\sum_c K_c = R_2$. Thus the gauge coupling $g_{a,b,c}$ associated to the $U(n_{a,b,c})$ is given by

$$\frac{1}{g^2_{a,b,c}} = \frac{I_a J_b K_c}{g_s \ell_s}. \tag{I.1}$$

Theta angles

The interaction which gives rise to the theta angle of the gauge theories in two dimensions can be seen to arise from the following interaction in the world-volume of the M-theory fivebrane

$$\int_{W_6} C^{(3)} \wedge dB^{(2)}. \tag{I.2}$$

It is well known that the fivebrane of M-theory wrapped on $\mathbf{S}^1$ is precisely the $D4$-brane of type IIA superstring theory. Thus the fivebrane will extend in the directions $x_0, x_1, x_2, x_4, x_6, x_{10}$ with x_{10} parametrizes $\mathbf{S}^1$. Interaction (I.2) turns out to be

$$\int_{(01246)\times\mathbf{S}^1} C^{(3)} \wedge dB^{(2)}, \tag{I.3}$$

with (01) the world-sheet coordinates where is defined the two-dimensional gauge theory. Under compactification on the circle $B^{(2)}$ field becomes the one-form A on the world-volume of the D4 branes. Eq. (I.3) yields

$$\int_{(01246)} C^{(3)} \wedge dA, \tag{I.4}$$

Under the further dimensional reduction of the D4 brane on 246 we obtain the two-dimensional field theory theta terms

$$\theta \int_{(01)} F, \tag{I.5}$$

where $\theta = \int_{(246)} C^{(3)}$.

There is another equivalent way to obtain the interaction (I.4), from the Chern-Simons contribution to the D-brane Lagrangian [33]

$$\int_{W_{p+1}} C \wedge \mathrm{Tr} e^{\mathcal{F}}, \tag{I.6}$$

where C is the sum over all RR fields of the relevant string theory and $\mathcal{F}$ is the field strength of the gauge fields on W_{p+1}. For D4 brane of type IIA string theory Chern-Simons Lagrangian looks like

$$\int_{W_5} \Big(C^{(1)} \mathrm{Tr}(\mathcal{F} \wedge \mathcal{F}) + C^{(3)} Tr \mathcal{F} \Big), \tag{I.7}$$

with $C^{(1)}$ and $C^{(3)}$ the IIA one- and three-forms. The last term reproduces Eq.(I.4).

The above analysis gives the correct answer only for the case of D4 branes wrapped on a $\mathbf{T}^3$, which gives two-dimensional gauge theories with sixteen supercharges. In more

general cases, the theta angle receives further contributions, which we analyze in the following.

Let $R_{a,b,c}$ denote the volume of a 246 box which is bounded by NS, NS$'$ and NS$''$ branes. We define Q_a to be the face contained in the a^{th} NS brane, Q_b the face contained in the b^{th} NS$'$ brane, and Q_c the face in the c^{th} NS$''$ brane. Let us consider the two-form fields $B^{(2)}$, $B'^{(2)}$ and $B''^{(2)}$ on the world-volume of NS, NS$'$ and NS$''$ branes. The theta angle for the group associated to the box (a, b, c) depends on the surface integrals along the boundary of $R_{a,b,c}$.

Let us first consider models with $(4, 4)$ supersymmetry in two dimensions. We can construct such models by considering $k' = k'' = 0$, and 246 compact. Each factor in the group arises from a box with the topology of a two-torus (along 2,4) times an interval (bounded by the positions of NS branes along 6). In such situation, the definition $\theta = \int_{R_{a,b,c}} C^{(3)}$ is not invariant under the gauge transformation of the three-form $C^{(3)} \to C^{(3)} + d\lambda$, with λ a two-form, since θ would pick a boundary term. This problem is solved by refining the definition as follows

$$\theta_a = \int_{R_{a,b,c}} C^{(3)} + \int_{Q_a} (B_a^{(2)} - B_{a-1}^{(2)}). \tag{I.8}$$

and allowing $B^{(2)}$ to transform as $B^{(2)} \to B^{(2)} + \lambda$.

In the general case in which the box $R_{a,b,c}$ is bounded by NS fivebranes in all three directions 246, the expression for the theta angle includes all the boundary terms.

$$\theta_{a,b,c} = \int_{R_{a,b,c}} C^{(3)} + \int_{Q_b} (B_b^{(2)} - B_{b-1}^{(2)}) + \int_{Q_a} (B_{a-1}^{\prime(2)} - B_a^{\prime(2)}) + \int_{Q_c} (B_c^{\prime\prime(2)} - B_{c-1}^{\prime\prime(2)}) \tag{I.9}$$

Now this expression is not invariant under the gauge transformations of the two-form fields, $B_a^{(2)} \to B_a^{(2)} + d\alpha_a$, etc, (here the α's are one-forms). Under this transformations, θ would pick up terms at the boundaries of the Q's, *i.e* intervals where two kinds of NS fivebranes intersect. This forces a further refinement by introducing integrals of appropriate one-form fields along those intervals. Finally, there should exist a contribution from the points of intersection of the three kinds of NS fivebranes, in order to ensure invariance under gauge transformations of the one-forms. We have not written these last contributions in the formula (I.9).

Fayet-Illiopoulos parameters

In two dimensions, we expect the gauge groups to contain $U(1)$ factors, and so the two-dimensional field theory contains Fayet-Iliopoulos parameters. These parameters will be associated to the positions of fivebranes in several directions. There are three contributions to the FI parameter of a given gauge group, one for each type of NS fivebrane bounding the corresponding box. One contribution comes from the position of the NS branes in x^7. For example, consider the case of k arbitrary and $k' = k'' = 1$. For each one of the k boxes we have a U(1)-factor and correspondingly a FI parameter. This is given by

$$r_a = (x^7)_{NS_a} - (x^7)_{NS_{a+1}} \equiv \delta(x^7)_a. \tag{I.10}$$

These FI parameters fulfill the condition $\sum_a^{k-1} r_a = 0$ required in order to have unbroken supersymmetry. The motivation for equation (I.10) is that the Higgs breaking triggered by the contribution of the FI to the D-term is precisely that obtained upon moving the NS brane in x^7.

This argument can be extended to the general case with arbitrary number of boxes. In this case FI parameters are given in terms of the differences of positions of NS branes in x^7, NS$'$ branes in x^5, and NS$''$ branes in x^3, as follows

$$r_{a,b,c} = (\delta x^7)_a + (\delta x^5)_b + (\delta x^3)_c. \tag{I.11}$$

They verify the condition $\sum_{a,b,c} r_{a,b,c} = 0$.

In general we expect that, as happened with the theta angles, the field theory parameters will receive contributions associated to the intersections of NS fivebranes. This was already observed in [10], and is expected to happen in our case in even a more complicated manner.

References

[1] A. Giveon and D. Kutasov, "Brane Dynamics and Gauge Theory", hep-th/9802067.

[2] A. Hanany, E. Witten, "Type IIB Superstrings, BPS Monopoles and Three-Dimensional Gauge Dynamics", Nucl. Phys. B492 (1997) 152, hep-th/9711230.

[3] I. Brunner, A. Karch "Branes and Six Dimensional Fixed Points", Phys. Lett. B409 (1997) 109-116, hep-th/9705022; "Branes and Orientifolds Versus Hanany-Witten in Six Dimensions", JHEP 03(1998)003, hep-th/9712143. A. Hanany, A. Zaffaroni, "Branes and Six-Dimensional Supersymmetric Theories", hep-th/9712145.

[4] N. Seiberg, "Non-trivial Fixed Points of the Renormalization Group in Six Dimensions", Phys. Lett. B390 (1997) 169, hep-th/9609161.

[5] K. Intriligator, "RG Fixed Points in Six Dimensions Via Branes at Orbifold Singularities", Nucl. Phys. B496 (1997) 177, hep-th/9702038. J. Blum, K. Intriligator, "Consistency Conditions for Branes at Orbifold Singularities", Nucl.Phys. B506 (1997) 223, hep-th/9705030; "New Phases of String Theory and 6d RG Fixed Points Via Branes at Orbifold Singularities", Nucl. Phys. B506 (1997) 199, hep-th/9705044

[6] A. Hanany and A. Zaffaroni, "On the Realization of Chiral Four-dimensional Gauge Theories Using Branes", JHEP 05(1998)001, hep-th/9801134.

[7] J. L. F. Barbón, "Rotated Branes and $N = 1$ Duality", Phys. Lett. B402 (1997) 59, hep-th/9703051.

[8] K. Landsteiner, E. López, D. A. Lowe, "Duality of Chiral $N = 1$ Supersymmetric Gauge Theories Via Branes", JHEP 02(1998)007, hep-th/9801002. I. Brunner, A. Hanany, A. Karch, D. Lüst, "Brane Dynamics and Chiral non-Chiral Transitions", hep-th/9801017. S. Elitzur, A. Giveon, D. Kutasov, D. Tsabar, "Branes, Orientifolds and Chiral Gauge Theories", hep-th/9801020.

[9] J. Lykken, E. Poppitz, S. P. Trivedi, "Chiral gauge theories from D-branes", Phys.Lett. B416(1998)286, hep-th/9708134; "M(ore) on chiral gauge theories from D-branes", hep-th/9712193.

[10] A. Hanany, M. Strassler and A.M. Uranga, "Finite Theories and Marginal Operators on the Brane", hep-th/9803086.

[11] L. Randall, Y. Shirman, R. von Unge, "Brane Boxes: Bending and the Beta Function", hep-th/9806092.

[12] A. Hanany and A.M. Uranga, "Brane Boxes and Branes on Singularities", hep-th/9805139.

[13] M.R. Douglas and G. Moore, "D-branes, Quivers and ALE Instantons, hep-th/9603167.

[14] C. V. Johnson, R. C. Myers, "Aspects of Type IIB Theory on ALE Spaces", Phys. Rev. D55 (1997) 6382, hep-th/9610140.

[15] K. Mohri, "D-branes and Quotients Singularities of Calabi-Yau Fourfolds", Nucl. Phys B521(1998)161, hep-th/9707012.

[16] M.R. Douglas, B.R. Greene and D.R. Morrison, "Orbifolds Resolution by D-branes", Nucl. Phys. B506(1997)84, hep-th/9704151.

[17] A. Lawrence, N. Nekrasov and C. Vafa, "On Conformal Field Theories in Four Dimensions", hep-th/9803015.

[18] L. E. Ibáñez, "A Chiral $D = 4$, $N = 1$ String Vacuum with a Finite Low Energy Effective Field Theory", hep-th/9802103.

[19] S. Kachru and E. Silverstein, "4d Conformal Field Theories and Strings on Orbifolds", Phys. Rev. Lett. 80(1998)4855, hep-th/9802183.

[20] M. Berkooz, M. R. Douglas, R. G. Leigh, "Branes Intersecting at Angles", Nucl.Phys. B480 (1996) 265.

[21] A. Hanany and K. Hori, "Branes and $N = 2$ Theories in Two Dimensions", Nucl. Phys. B513(1998)119, hep-th/9707192.

[22] J. H. Brodie, "Two Dimensional Mirror Symmetry From M-theory", Nucl.Phys. B517 (1998) 36, hep-th/9709228. M. Alishahiha, "N=(4,4) 2D Supersymmetric Gauge Theory and Brane Configuration", Phys. Lett. B420 (1998) 51, hep-th/9710020; "On the Brane Configuration of $N = (4, 4)$ 2D Supersymmetric Gauge Theory", hep-th/9802151.

[23] W. Lerche, "Fayet-Iliopoulos Potentials from Four-Folds", JHEP 11(1998)004, hep-th/9709146.

[24] E. Witten, "Phases of $N = 2$ Theories in Two Dimensions", Nucl. Phys. B403(1993)159, hep-th/9301042.

[25] J. Distler, "Notes on (0,2) Superconformal Field Theories", hep-th/9502012.

[26] E. Witten. "Solutions of Four-dimensional Field Theories Via M Theory", Nucl. Phys. B500 (1997) 3.

[27] E. Witten, "Sigma Models and the ADHM Construction of Instantons", J. Geom. Phys. 15 (1995) 215, hep-th/9410052.

[28] K. Dasgupta, S. Mukhi, "A Note on Low-Dimensional String Compactifications", Phys. Lett. B398 (1997) 285, hep-th/9612188.

[29] T. Banks, N. Seiberg, E. Silverstein, "Zero and One-dimensional Probes with $N = 8$ Supersymmetry", Phys. Lett. B401 (1997) 30, hep-th/9703052.

[30] P. S. Aspinwall, B. R. Greene, D. R. Morrison, "Calabi-Yau Moduli Space, Mirror Manifolds and Spacetime Topology Change in String Theory", Nucl.Phys. B416(1994)414.

[31] T. Muto, "D-branes on Orbifolds and Topology Change", hep-th/9711090; B. R. Greene, "D-Brane Topology Changing Transitions", hep-th/9711124; S. Mukhopadhyay, K. Ray, "Conifolds from D-branes", Phys. Lett. B423(1998)62.

[32] S.P. de Alwis, "Coupling of Branes and Normalization of Effective Actions in String/M-theory", Phys. Rev. D56(1997)7963, hep-th/9705139.

[33] M.R. Douglas, "Branes within Branes", hep-th/9512077.